


HVP CONTRIBUTION TO $(g - 2)_\mu$: STATUS OF THE MAINZ CALCULATION

M. CÈ, A. GÉRARDIN, G. VON HIPPEL, R. J. HUDSPITH, S. KUBERSKI, H. B. MEYER,
K. MIURA, D. MOHLER, K. OTTNAD, S. PAUL, A. RISCH, T. SAN JOSÉ, H. WITTIG

FIFTH PLENARY WORKSHOP OF THE MUON $g - 2$ THEORY INITIATIVE
SEPTEMBER 5, 2022

HADRONIC VACUUM POLARIZATION CONTRIBUTION TO THE MUON $g - 2$



← Status for a_μ^{hvp} [2203.15810, Colangelo et al.]


- Prediction in [2002.12347, BMWc] deviates **significantly** from data-driven results.
- High-precision lattice calculations needed. Major challenges:
 - ▶ Cutoff effects at short distances t
 - ▶ Exponential deterioration of signal-to-noise ratio at large t (with traditional Monte Carlo methods)

- Short term: Focus on benchmark quantities to compare among collaborations. Time windows in the Time Momentum Representation [1801.07224, Blum et al.]
- Long term: Improve overall precision of a_μ^{hvp} .

EUCLIDEAN TIME WINDOWS IN THE TMR: ISOVECTOR CHANNEL

Time-momentum representation [1107.4388, Bernecker and Meyer]:

$$(a_\mu^{\text{hvp}})^i := \left(\frac{\alpha}{\pi}\right)^2 \int_0^\infty dt \tilde{K}(t) G(t) W^i(t; t_0; t_1)$$



■ Current-current correlator:

$$G(t) = -\frac{a^3}{3} \sum_{k=1}^3 \sum_{\vec{x}} \langle j_k^{\text{em}}(t, \vec{x}) j_k^{\text{em}}(0) \rangle$$

■ Time windows [1801.07224, Blum et al.]:

$$W^{\text{SD}}(t; t_0; t_1) = [1 - \Theta(t, t_0, \Delta)]$$

$$W^{\text{ID}}(t; t_0; t_1) = [\Theta(t, t_0, \Delta) - \Theta(t, t_1, \Delta)]$$


$$W^{\text{LD}}(t; t_0; t_1) = \Theta(t, t_0, \Delta)$$

where

$$\Theta(t, t', \Delta) := \frac{1}{2} (1 + \tanh[(t - t')/\Delta])$$

$$t_0 = 0.4 \text{ fm}, t_1 = 1.0 \text{ fm}, \Delta = 0.15 \text{ fm}.$$

2 + 1 FLAVOR CLS ENSEMBLES



- $O(a)$ improved Wilson-clover fermions.
- Six values of $a \in [0.039, 0.099]$ fm, a factor of 6.4 in a^2 .
- Open boundary conditions in temporal direction.
- $m_\pi \in [129, 422]$ MeV
- $a\text{Tr}[M_q] = 2am_l + am_s = \text{const.}$

Scale: Either use $\sqrt{t_0^{\text{phys}}} = 0.1443(15)$ fm [2112.06696, Straßberger et al.] or express dimensionfull quantities in terms of af_π [1103.4818, Xu et al.][1904.03120, Gérardin et al.]

→ new $N_f = 2 + 1$ result by RQCD: $\sqrt{t_0^{\text{phys}}} = 0.1449_{(9)}^{(7)}$ fm may be used in the future.

- Work in isospin decomposition of the electromagnetic current

$$j_\mu^{\text{em}} = \frac{2}{3}\bar{u}\gamma_\mu u - \frac{1}{3}\bar{d}\gamma_\mu d - \frac{1}{3}\bar{s}\gamma_\mu s + \frac{2}{3}\bar{c}\gamma_\mu c + \dots = j_\mu^{I=1} + j_\mu^{I=0} + \frac{2}{3}\bar{c}\gamma_\mu c + \dots, ,$$

$$\text{Isovector: } j_\mu^{I=1} = \frac{1}{2}(\bar{u}\gamma_\mu u - \bar{d}\gamma_\mu d), \quad \text{Isoscalar: } j_\mu^{I=0} = \frac{1}{6}(\bar{u}\gamma_\mu u + \bar{d}\gamma_\mu d - 2\bar{s}\gamma_\mu s)$$

- Two discretizations of the vector current: local and conserved

$$J_\mu^{(\text{L}),a}(x) = \bar{\psi}(x)\gamma_\mu \frac{\lambda^a}{2}\psi(x),$$

$$J_\mu^{(\text{C}),a}(x) = \frac{1}{2} \left(\bar{\psi}(x + a\hat{\mu})(1 + \gamma_\mu)U_\mu^\dagger(x) \frac{\lambda^a}{2}\psi(x) - \bar{\psi}(x)(1 - \gamma_\mu)U_\mu(x) \frac{\lambda^a}{2}\psi(x + a\hat{\mu}) \right),$$

$O(a)$ IMPROVED VECTOR CURRENTS

- Improved vector currents are given by

$$J_\mu^{(\alpha),a,I}(x) = J_\mu^{(\alpha),a}(x) + ac_V^{(\alpha)}(g_0) \tilde{\partial}_\nu \Sigma_{\mu\nu}^a(x), \quad \text{with } \alpha \in L, C$$

- Renormalization and mass-dependent improvement of local currents via

$$J_\mu^{(L),3,R}(x) = Z_V [1 + 3\bar{b}_V am_q^{\text{av}} + b_V am_{q,l}] J_\mu^{(L),3,I}(x),$$

$$J_\mu^{(L),8,R}(x) = Z_V \left[1 + 3\bar{b}_V am_q^{\text{av}} + \frac{b_V}{3} a(m_{q,l} + 2m_{q,s}) \right] J_\mu^{(L),8,I}(x) \\ + Z_V \left(\frac{1}{3} b_V + f_V \right) \frac{2}{\sqrt{3}} a(m_{q,l} - m_{q,s}) J_\mu^{(L),0,I}(x),$$

- Two independent non-perturbative determinations of Z_V , c_V^L , c_V^C , b_V , \bar{b}_V :

Set 1: Large-volume, CLS ensembles [1811.08209, Gérardin et al.]

Set 2: Small volume, Schrödinger functional [2010.09539, ALPHA],[1805.07401, Fritzsche]

differ by higher order cutoff effects. f_V is of $O(g_0^6)$ and unknown.


FINITE-SIZE EFFECTS

- Finite-size corrections applied to the isovector correlator.
- Correction for $t < \frac{(m_\pi L/4)^2}{m_\pi}$: Hansen-Patella method [1904.10010][2004.03935]
 - ▶ Expansion in the pion winding number.
 - ▶ Using monopole parametrization of the electromagnetic pion form factor.
- Large distances: MLL [1105.1892, Meyer] [hep-lat/0003023, Lellouch and Lüscher]:
 - ▶ Compute difference between finite and infinite-volume isovector correlator
 - ▶ Based on the time-like pion form factor.
 - ▶ Applied at large Euclidean distances \rightarrow less relevant for short and intermediate distance windows.
- This is the **only correction** applied to the lattice data!
Of similar size as statistical uncertainty for $a_\mu^{\text{win}} \equiv (a_\mu^{\text{hvp}})^{\text{ID}}$.

THE INTERMEDIATE-DISTANCE WINDOW


[2206.06582, Cè et al.]

CONTINUUM EXTRAPOLATION AT $SU(3)_f$ SYMMETRIC POINT



- Two sets of equally valid improvement coefficients.
- No cutoff effects of $O(a^3)$ resolved for Set 1.
- Independent extrapolations compatible in the continuum \rightarrow strong cross-check of our extrapolations.
- No sign of modification $a^2 \rightarrow (\alpha_s(1/a^2))^{\hat{\Gamma}} a^2$
[1912.08498, Husung et al.]


CHIRAL EXTRAPOLATION OF ISOVECTOR CONTRIBUTION




- f_π rescaling, local-local current and Set 1.
- Curvature in $\tilde{y} = \frac{m_\pi^2}{8\pi f_\pi^2}$ is needed to describe the data.
- Singular fit ansatz favored, also found in [2110.05493, Colangelo et al.]
- Variation in the chiral extrapolation does not change the result significantly.

$$a_\mu^{\text{win}}(\tilde{y}) = \gamma_1 (\tilde{y} - \tilde{y}^{\text{exp}}) + \gamma_2 (f(\tilde{y}) - f(\tilde{y}^{\text{exp}})) , \quad f(\tilde{y}) \in \{0; \log(\tilde{y}); \tilde{y}^2; 1/\tilde{y}; \tilde{y} \log(\tilde{y})\}$$


MODEL AVERAGES: ISOVECTOR CONTRIBUTION



- Eight combinations of discretization and improvement procedures.
- Model averages in each category to determine systematic uncertainty from choice of fit model. [\[2008.01069, Jay and Neil\]](#)
- Final result by combining L and C of Set 1.




COMPARISON WITH LATTICE RESULTS FOR $a_\mu^{\text{win,iso}}$



$$a_\mu^{\text{win,iso}} = a_\mu^{\text{win,I1}} + a_\mu^{\text{win,I0}} + a_\mu^{\text{win,c}} = (236.60 \pm 0.79_{\text{stat}} \pm 1.13_{\text{syst}} \pm 0.05_{\text{Q}}) \times 10^{-10}$$

- Tension with EMTC 21 and RBC/UKQCD 18 estimates for $a_\mu^{\text{win,iso}}$ mainly from light quark contribution. EMTC 22 and BMW 20 consistent with our result.

ISOSPIN BREAKING EFFECTS IN a_μ^{win}



- QED_L-action [0804.2044, Hayakawa and Uno] for IR regularisation, Coulomb gauge.


- Reweighting based on perturbative expansion [1303.4896, de Divitiis et al.] in $\Delta\varepsilon = \varepsilon - \varepsilon^{(0)} = (\Delta m_u, \Delta m_d, \Delta m_s, \Delta\beta = 0, e^2)$

- More details in [Andreas Risch's contribution](#).
- IB in scale setting [2112.08262, Segner et al.] and QED-FV effects to be considered.
- Uncertainty on relative correction 0.3(1)% doubled in final result for a_μ^{win} .

COMPARISON WITH RESULTS FOR a_μ^{win}

- Isospin-breaking correction $+(0.70 \pm 0.47) \times 10^{-10}$ included:

$$a_\mu^{\text{win}} = (237.30 \pm 0.79_{\text{stat}} \pm 1.13_{\text{syst}} \pm 0.05_{\text{Q}} \pm 0.47_{\text{IB}}) \times 10^{-10}$$



- 3.9σ tension with data-driven estimate in [2205.12963, Colangelo et al.].
- Genuine difference between lattice and data-driven results?



INTERMEDIATE WINDOW: COVARIANT-COORDINATE SPACE METHOD

- The covariant-coordinate space (CCS) method [1706.01139, Meyer] offers an alternative to the TMR to compute a_μ^{hvp} and a_μ^{win} :

$$a_\mu^{\text{win}} = \int d^4x H_{\mu\nu}(x) G_{\mu\nu}(x),$$
$$H_{\mu\nu} = -\delta_{\mu\nu} \mathcal{H}_1(|x|) + \frac{x_\mu x_\nu}{|x|^2} \mathcal{H}_2(|x|), \quad G_{\mu\nu}(x) = \langle j_\mu(x) j_\nu(0) \rangle$$


- Freedom of choice in kernel functions $H_{\mu\nu}$ [1811.08669, Cè et al.].
- Based on integration in four-dimensional sphere. Especially suited for very large lattices.
- Computation of a_μ^{win} on CLS lattices including new treatment of finite volume effects by Chao, Meyer, Parrino (Julian Parrino's talk at Lattice22) as cross-check of TMR results.

INTERMEDIATE WINDOW: COVARIANT-COORDINATE SPACE METHOD



- Continuum limit of CCS data and comparison with TMR at $m_\pi \sim 346-353$ MeV, presented in [Julian Parrino's talk at Lattice22](#).

INTERMEDIATE WINDOW: COVARIANT-COORDINATE SPACE METHOD



- CCS results shifted to the physical pion mass using information from the TMR method (10% correction).

- Continuum limit:

$$(a_\mu^{\text{win,II}})_{\text{II}} = 186.11(1.42) \cdot 10^{-10}$$

$$(a_\mu^{\text{win,II}})_{\text{cl}} = 185.01(1.40) \cdot 10^{-10}$$

to be compared with the TMR result

$$a_\mu^{\text{win,II}} = 186.30(0.75)(1.08) \cdot 10^{-10}$$

→ excellent agreement between both methods.


Presented in [Julian Parrino's talk at Lattice22](#).

THE SHORT-DISTANCE WINDOW

THE SHORT-DISTANCE WINDOW

- Short-distance cutoff effects are dominant source of uncertainty in $(a_\mu^{\text{hvp}})^{\text{SD}}$.
 - Log-enhanced cutoff effects are present at very short distances of the TMR integral [0807.1120, Della Morte et al.][2106.15293, Cè et al.] [Rainer Sommer's talk at Lattice22] → also present in a_μ^{hvp} !
-
- Tree-level improvement may help to reduce cutoff effects, as used for $(a_\mu^{\text{hvp}})^{\text{SD}}$ in [2206.15084, Alexandrou et al.].
 - Use of perturbation theory at $O(\alpha_s^4)$ at very short distances (already suggested in [1107.4388, Bernecker and Meyer]) removes logarithmic contribution.

THE REGULATED SHORT-DISTANCE WINDOW



- Regulate the short distance part [Rainer Sommer's talk at Lattice22]:

$$\int_0^\infty dt F(t) = \int_0^\infty dt [1 - \chi(t)] F(t) + \lim_{a \rightarrow 0} a \sum_0^\infty \chi(t) F(t).$$

Combine **perturbation theory at short distances** with the **continuum lattice result**.


- Choose sufficiently smooth and short ranged regulator, e.g.,

$$\chi(t) = \theta(t - u_0) \left(1 - \cos \left[\frac{(t - u_0)\pi}{(2\delta)} \right]^2 \theta(u_0 + \delta - t) \right)$$

with the Heaviside step function $\theta(t)$ and $u_0 = \delta = 0.075$ fm.


THE REGULATED SHORT-DISTANCE WINDOW

- Test regulators, improvement schemes and discretization prescriptions.
- Continuum extrapolation of regulated short-distance window $(a_\mu^{\text{HVP,II}})^{\text{SD,P}}$ at the $\text{SU}(3)_f$ -symmetric point.




- No log-enhanced cutoff effects expected.
- Tree-level improvement:
 - ▶ Based on massless, free theory.
 - ▶ Reduces cutoff effects at $a = 0.1$ fm from 18% to 6%.
- Different data sets offer insight in systematic uncertainties.


NOISE REDUCTION IN THE LONG-DISTANCE TAIL



- 2.2% uncertainty: Dominated by statistical uncertainties of light quark contribution.
- Variance reduction is needed to reach sub-percent precision (2.2% statistical uncertainty at physical pion mass).
- Computation of isospin breaking effects ongoing: [Andreas Risch's contribution](#).



VARIANCE REDUCTION: SPECTROSCOPY




- Dedicated spectroscopy analysis of finite-volume energies and amplitudes to reconstruct the tail of the isovector correlation function [1808.05007, Andersen et al.][1904.03120, Gérardin et al.].
- Analysis at close-to-physical masses ongoing [2112.07385, Paul et al.], up to 9 energy levels resolved.

- Spectral decomposition of the vector correlator:

$$G_l(t) = \sum_n |A_n|^2 e^{-E_n t}, \quad E_n = 2\sqrt{m_\pi^2 + k^2}$$


- Computation of the pion transition form factor to correct for finite-size effects with less model dependence ongoing.

VARIANCE REDUCTION: SPECTROSCOPY




- Reconstruction of the light-connected TMR correlator at long distances at close-to-physical pion mass, presented in [Srijit Paul's talk at Lattice22](#).

VARIANCE REDUCTION: SPECTROSCOPY



- Reconstruction of the light-connected TMR correlator at long distances at close-to-physical pion mass, presented in [Srijit Paul's talk at Lattice22](#).


VARIANCE REDUCTION: LOW MODE AVERAGING



- Low Mode Averaging (LMA) [[hep-lat/0106016](#), Neff et al.][[hep-lat/0402002](#), Giusti et al.] to reduce the variance of the isovector contribution.
- Light-connected contribution to a_μ^{hvp} close-to-physical pion mass in a $12.4 \text{ fm} \times (6.2 \text{ fm})^3$ box at $a = 0.064 \text{ fm}$.


- 800 eigenmodes of the even-odd preconditioned Dirac-Wilson operator $\gamma_5 \hat{D}$.
- All-to-all evaluation of low eigenmodes dominates for $t > 1.5 \text{ fm}$.

VARIANCE REDUCTION: LOW MODE AVERAGING



- Comparison of stochastic evaluation and LMA at similar statistics.
- Ongoing calculation: Sub-percent precision reached.
- LMA also applied on less challenging ensembles.

VARIANCE REDUCTION: COMPARISON



- Comparison of statistical uncertainties of the TMR correlator based on stochastic evaluation, reconstruction and LMA.
- Significantly less noise in LMA correlator.
- Exponential noise reduction in reconstructed correlator.

Intermediate window

- We observe tension with data-driven estimates for a_μ^{win} .
- Systematic effects from continuum extrapolation seem to be under control:
 - ▶ Non-perturbative $O(a)$ improvement
 - ▶ 6 resolutions < 0.1 fm with $a_{\text{max}}^2/a_{\text{min}}^2 > 6$.
 - ▶ Two discretizations of the vector current, two sets of improvement procedures.
 - ▶ So-far no sign of logarithmic corrections to a^2 scaling.
- Uncertainties from chiral extrapolation and finite-volume correction are subleading.

- Similar tension found for $\Delta\alpha_{\text{had}}(Q^2) \rightarrow$ [Marco Cè's talk](#).
- Investigation of other windows might help to clarify the situation.
- Short-distance window:
 - ▶ Cutoff effects from short-distance singularities need proper treatment [[0807.1120, Della Morte et al.](#)][[2106.15293, Cè et al.](#)] [[Rainer Sommer's talk](#)].
 - ▶ Systematic uncertainties will dominate and need to be properly estimated.
- Sub-percent precision on a_{μ}^{hvp} needs reduction of our statistical uncertainties.
- Spectroscopy and variance reduction techniques will help to improve our calculation significantly close to physical pion mass.