

Dispersive analysis of the isospin-breaking corrections to
 $e^+e^- \rightarrow \pi^+\pi^-$ and $\pi^+\pi^- \rightarrow \pi^+\pi^-$

J. Ruiz de Elvira

Departamento de Física Teórica, Universidad Complutense de Madrid

Fifth Plenary Workshop of the Muon g-2 Theory Initiative
September 6th 2022



Introduction

Interference: RC to the forward-backward asymmetry in $e^+e^- \rightarrow \pi^+\pi^-$

Isospin-breaking corrections for $\pi\pi$ scattering

Dispersive approach to FSR in $e^+e^- \rightarrow \pi^+\pi^-$

Summary / Outlook

Work in collaboration with

Gilberto Colangelo, Martin Hoferichter and Joachim Monnard

Introduction

Interference: RC to the forward-backward asymmetry in $e^+e^- \rightarrow \pi^+\pi^-$

Isospin-breaking corrections for $\pi\pi$ scattering

Dispersive approach to FSR in $e^+e^- \rightarrow \pi^+\pi^-$


Summary / Outlook

| Contribution | Value $\times 10^{11}$ |
|--|------------------------|
| QED | 116 584 718.931(104) |
| Electroweak | 153.6(1.0) |
| HVP ($e^+ e^-$, LO + NLO + NNLO) | 6845(40) |
| HLbL (phenomenology + lattice + NLO) | 92(18) |
| Total SM Value | 116 591 810(43) |
| Experiment | 116 592 061(41) |
| Difference: $\Delta a_\mu := a_\mu^{\text{exp}} - a_\mu^{\text{SM}}$ | 251(59) |

- HVP **dominant** source of theory **uncertainty**
relative size of $\Delta\text{HVP} \sim 0.6\%$
- 2π channel provides **70%** of the HVP contribution
 \hookrightarrow RC in $e^+ e^- \rightarrow \pi^+ \pi^-$ must be **under control**
- RC evaluation based on **models** so far
 \hookrightarrow a **dispersive** approach could lead to **model-independent** results


[Talk from P. Stoffer]

- Initial State Radiation:



can be calculated in QED in terms of $F_\pi^V(s)$

- Interference terms




- require hadronic matrix elements **beyond $F_\pi^V(s)$**
- so far **estimated** using sQED+ $F_\pi^V(s)$ or (generalized) VMD **models**

[Arbuzov, Kopylova, Seilkhanova (2020), Ignatov, Lee (2022)]

- **pion-pole** contribution analyzed **dispersively**, this talk

[Colangelo, Hoferichter, Monnard, JRE (2022)]

- Final State Radiation:




- also requires hadronic matrix elements **beyond** $F_\pi^V(s)$

- known in ChPT to one loop

[Kubis, Meißner (2001)]

↪ **dispersive determination** this talk

Dispersive approach to FSR



- Neglecting intermediate states beyond 2π , unitarity reads

$$\begin{aligned}
 \text{Im } F_V^{\pi, \alpha}(s) &= \int d\phi_2 F_V^{\pi}(s) \times T_{\pi\pi}^{\alpha}(s, t)^* \\
 &+ \int d\phi_2 F_V^{\pi, \alpha}(s) \times T_{\pi\pi}(s, t)^* \\
 &+ \int d\phi_3 F_V^{\pi, \gamma}(s, t) \times T_{\pi\pi}^{\gamma}(s, t')^*
 \end{aligned}$$

- Need $T_{\pi\pi}^{\alpha}$ as well as $F_{\pi}^{V, \gamma}$ and $T_{\pi\pi}^{\gamma}$ as input

↔ dispersive approach to RC to $\pi\pi$ scattering

- The DR for $F_{\pi}^{V, \alpha}(s)$ takes the form of an integral equation

Introduction


Interference: RC to the forward-backward asymmetry in $e^+e^- \rightarrow \pi^+\pi^-$

Isospin-breaking corrections for $\pi\pi$ scattering

Dispersive approach to FSR in $e^+e^- \rightarrow \pi^+\pi^-$

Summary / Outlook

- Interference terms: **pion-pole** contribution



- do not contribute to the total cross section
can be **tested** in the **forward-backward** asymmetry

[CMD-3 results, talk from Ivan Logashenko]

$$A_{\text{FB}}(z) = \frac{\frac{d\sigma}{dz}(z) - \frac{d\sigma}{dz}(-z)}{\frac{d\sigma}{dz}(z) + \frac{d\sigma}{dz}(-z)}, \quad z = \cos \theta,$$

non-vanishing from **RC**, **C-odd** terms

- Box diagram** contributes together to **ISR-FSR** soft radiation

$$\left. \frac{d\sigma}{dz} \right|_{\text{C-odd soft}} = \frac{d\sigma_0}{dz} \left[\delta_{\text{soft}}(m_\gamma^2, \Delta) + \delta_{\text{virt}}(m_\gamma^2) \right]$$

Forward-backward asymmetry in $e^+e^- \rightarrow \pi^+\pi^-$


- δ_{soft} computed analytically in QED

$$\delta_{\text{soft}} = \frac{2\alpha}{\pi} \left\{ \log \frac{m_\gamma^2}{4\Delta^2} \log \frac{1 + \beta z}{1 - \beta z} + \log(1 - \beta^2) \log \frac{1 + \beta z}{1 - \beta z} + \dots \right\},$$

[Arbuzov et al. (2020), Ignatov, Lee (2022), Colangelo, Hoferichter, Monnard, JRE (2022)]

- δ_{virt} computed dispersively

▷ start from a fixed-s dispersion relation



↔ for scalar particles D_0 function


▷ for real pions: dispersive representation of $F_\pi^V(s)$

$$\frac{F_\pi^V(s)}{s} = \frac{1}{s} + \frac{1}{\pi} \int_{4m_\pi^2}^{\infty} ds' \frac{\text{Im} F_\pi^V(s')}{s'(s' - s)} \rightarrow \frac{1}{s - m_\gamma^2} - \frac{1}{\pi} \int_{4m_\pi^2}^{\infty} ds' \frac{\text{Im} F_\pi^V(s')}{s'} \frac{1}{s - s'}$$

↔ the VFF corrections can be interpreted as a propagator

Forward-backward asymmetry in $e^+e^- \rightarrow \pi^+\pi^-$: results

- δ_{virt} decomposed in **pole-pole**, **pole-disp** and **disp-disp** contributions
- **pole-pole** and **pole-disp** IR divergent
 - ↔ cancel against the real emission



- disp-pole term dominates: **infrared enhancement**
- significant **corrections** beyond sQED
- similar results from GVMD models

[Colangelo, Hoferichter, Monnard, JRE (2022)]

[Ignatov, Lee (2022)]

Introduction

Interference: RC to the forward-backward asymmetry in $e^+e^- \rightarrow \pi^+\pi^-$

Isospin-breaking corrections for $\pi\pi$ scattering

Dispersive approach to FSR in $e^+e^- \rightarrow \pi^+\pi^-$

Summary / Outlook

- Starting point: Roy-equation solution for $\pi\pi$ scattering below $s_1 \sim 1$ GeV

[Ananthanarayan, Colangelo, Gasser, Leutwyler (2001), Garcia-Martin, Kaminski, Pelaez, JRE (2011)]

- $\pi\pi$ invariant amplitude

$$A(s, t, u) = A(s, t, u)_{SP} + A(s, t, u)_d$$

- A_{SP} contribution of S and P waves below s_1

$$A(s, t, u)_{SP} = 32\pi \left\{ \frac{1}{3} W^0(s) + \frac{3}{2}(s-u)W^1(t) + \frac{1}{2}W^2(t) + (t \leftrightarrow u) \right\}$$

- ▷ $W^l(s)$ only RHC, DR in terms of the S and P partial waves t_l^j

$$W^0(s) = \frac{a_0^0 s}{4M_\pi^2} + \frac{s(s-4M_\pi^2)}{\pi} \int_{4M_\pi^2}^{s_1} ds' \frac{\text{Im } t_0^0(s')}{s'(s'-4M_\pi^2)(s'-s)},$$

- A_d is the “background amplitude”, higher partial waves and higher energies
↔ for $s < s_1$ small and smooth, polynomial
- Construct isospin amplitudes T^0 , T^1 and T^2

- Three different **isospin-breaking** effects
 1. **strong** isospin breaking: effects proportional $(m_u - m_d)$
 2. effects proportional to $M_{\pi^+} - M_{\pi^0}$
 3. further **photon exchanges**
- Each of them can be **considered separately** from the other two

- At **low energies** chiral symmetry imposes $\mathcal{O}((m_u - m_d)^2)$

↪ **small** shift in M_{π^0}

[Gasser, Leutwyler (84)]

- Higher energies, generate $\pi^0 - \eta$ and $\rho - \omega$ mixing

- $\pi^0 - \eta$ not relevant for F_π^V : can be estimated phenomenologically

rescattering effects can be estimated from $\eta \rightarrow 3\pi$

[Colangelo, Lanz, Leutwyler, Passemar (2018)]

- $\rho - \omega$ mixing contribution allows for a high-precision description of F_π^V

[Colangelo, Hoferichter, Kubis, Stoffer (2022)]

1. ω meson described with a **narrow-width** approximation

2. $\rho - \omega$ **interference** through a single parameters ϵ_ω

3. ρ and ω coupling to **radiative channels** induces a **non-negligible phase**

- First, switch from the **isospin** to the **charge basis**

$$\Leftrightarrow T^0, T^1, T^2 \Rightarrow T^c, T^n, T^x$$

$$T^c := T(\pi^+\pi^- \rightarrow \pi^+\pi^-), \quad T^x := T(\pi^+\pi^- \rightarrow \pi^0\pi^0), \quad T^n := T(\pi^0\pi^0 \rightarrow \pi^0\pi^0)$$

- Adapt unitarity relation

$$\text{Im}t_{n,s}(s) = \sigma_0(s)|t_{n,s}(s)|^2 + 2\sigma(s)|t_{x,s}(s)|^2$$

$$\text{Im}t_{x,s}(s) = \sigma_0(s)t_{n,s}(s)t_{x,s}^*(s) + 2\sigma(s)t_{x,s}(s)t_{c,s}^*(s)$$

$$\text{Im}t_{c,s}(s) = \sigma_0(s)|t_{x,s}(s)|^2 + 2\sigma(s)|t_{c,s}(s)|^2$$

where

$$\sigma(s) = \sqrt{1 - \frac{4M_{\pi^+}^2}{s}}, \quad \sigma_0(s) = \sqrt{1 - \frac{4M_{\pi^0}^2}{s}}$$

\Leftrightarrow encode the effect of $M_{\pi^+} - M_{\pi^0}$

- Assume that the *input* above s_1 does **not change** for $M_{\pi^+}^2 - M_{\pi^0}^2 \neq 0$
- Concentrate in T_{SP} , S and P waves below ~ 1 GeV
- Express W^I in terms of the imaginary parts of the **physical channels**

$$T_{SP}^n(s, t, u) = 32\pi \left(W_{n,S}^{00}(s) + W_{n,S}^{+-}(s) + (s \leftrightarrow t) + (s \leftrightarrow u) \right)$$


where

$$W_{n,S}^{00}(s) = \frac{a_n^{00} s}{4M_{\pi^0}^2} + \frac{s(s - 4M_{\pi^0}^2)}{\pi} \int_{4M_{\pi^0}^2}^{s_1} ds' \frac{\text{Im}t_{n,S}^{00}(s')}{s'(s' - 4M_{\pi^0}^2)(s' - s)}$$

$$W_{n,S}^{+-}(s) = \frac{s(s - 4M_{\pi^0}^2)}{\pi} \int_{4M_{\pi^0}^2}^{s_1} ds' \frac{\text{Im}t_{n,S}^{+-}(s')}{s'(s' - 4M_{\pi^0}^2)(s' - s)}$$

- similar for the other channels

- 1 Starting point: take the isospin limit Roy-equation solution T_0^C, T_0^X, T_0^n
 - 2 Reevaluate the dispersive integrals with the shifted threshold
 - 3 Iterate the procedure until convergence
- Preliminary results:



- The effect on $F_{\pi}^V(s)$ is small
($\pi^0\pi^0$ only appears in the t-channel of the $\pi\pi$ amplitude in the unitarity relation)

Colangelo, Monnard, JRE (preliminary)


Roy equations and photon-exchange effects

- Photon-exchange diagrams are **not** included in Roy equations
- Modify Roy-equation solutions (T_0^i) to include $\mathcal{O}(\alpha)$ effects
- We start with the **Born term**

$$T_B(t, s, u) := \begin{array}{c} \pi^- \text{---} \bullet \text{---} \pi^- \\ | \\ \text{spring} \\ | \\ \pi^+ \text{---} \bullet \text{---} \pi^+ \end{array} = 4\pi\alpha \frac{s-u}{t} F_\pi^V(t)^2$$

contribution to $T_B^C(s, t, u) = T_B^C(t, s, u) + T^C(s, t, u)$


- Adding T_B^C to T^C affects **unitarity relations** for all amplitudes



↪ we are generating further $\mathcal{O}(\alpha)$ corrections: **iterative procedure**

Roy equations and photon-exchange effects: first iteration

- Remark: through this **procedure** we are not generating **box diagrams**



- Compute them through **double-spectral representation**

$$T_D^C(s, t, u) := \text{diagram 1} + \text{diagram 2} + \text{diagram 3} + \text{flipped diags.}$$

$$T_D^X(s, t, u) := \text{diagram}$$

- Include them as **starting point** for **further iterations**


$$T^C(s, t, u) = T_0^C(s, t, u) + T_B^C(s, t, u) + T_D^C(s, t, u)$$

$$T^X(s, t, u) = T_0^X(s, t, u) + T_D^X(s, t, u)$$

$$T^n(s, t, u) = T_0^n(s, t, u)$$

Roy equations and photon-exchange effects: further iterations

- For the **second iteration** we have the diagrams



- they have to be **cut** in **all possible ways**:
↪ contributions from subamplitudes with **real photons**: more later

- After N -iterations:

$$T^C(s, t, u) = T_0^C(s, t, u) + T_B^C(s, t, u) + T_D^C(s, t, u) + \sum_{k=2}^N R_k^C(s, t, u)$$

$$T^X(s, t, u) = T_0^X(s, t, u) + T_D^X(s, t, u) + \sum_{k=2}^N R_k^X(s, t, u)$$

$$T^n(s, t, u) = T_0^n(s, t, u) + \sum_{k=2}^N R_k^n(s, t, u)$$

- each iteration k is $\mathcal{O}(p^{2k})$ in the **chiral** expansion

- The evaluation of R_{k+1}^i , with $k \geq 1$ is done as follows:
 1. project the R_k^i amplitudes onto partial waves
 2. insert these into the **unitarity relations** combined with the projections of T_0^i
 3. add the contribution of subdiagrams with **real photons**
 4. **solve** the corresponding **dispersion relation**

- Subtraction constants can be fixed by matching to ChPT
 - ▷ ChPT $\pi\pi$ amplitude with RC known to one loop [Knecht, Urech (1997), Knecht, Nehme (2002)]

- Work **in progress**: **preliminary** results [J. Monnard thesis, \(2021\)](#)

Introduction

Interference: RC to the forward-backward asymmetry in $e^+e^- \rightarrow \pi^+\pi^-$

Isospin-breaking corrections for $\pi\pi$ scattering

Dispersive approach to FSR in $e^+e^- \rightarrow \pi^+\pi^-$

Summary / Outlook

$$\begin{aligned} \text{Im } F_V^{\pi,\alpha}(s) &= \int d\phi_2 F_V^{\pi}(s) \times T_{\pi\pi}^{\alpha}(s, t)^* \\ &+ \int d\phi_2 F_V^{\pi,\alpha}(s) \times T_{\pi\pi}(s, t)^* \\ &+ \int d\phi_3 F_V^{\pi,\gamma}(s, t) \times T_{\pi\pi}^{\gamma}(s, t')^* \end{aligned}$$

- After this long digression we have obtained **preliminary** results for $T_{\pi\pi}^{\alpha}$
- For $F_V^{\pi,\gamma}(s, t)$ and $T_{\pi\pi}^{\gamma}(s, t')$




- pion-pole contribution + $\gamma\gamma \rightarrow \pi\pi$ input


↪ all subamplitudes known: $F_V^{\pi,\gamma}(s, t)$ and $T_{\pi\pi}^{\gamma}(s, t')$ computed

Evaluation of $F_{\pi}^{V,\alpha}$

- Work in progress:
 1. $M_{\pi}^{+} - M_{\pi}^0$ effects missing
 2. controlled matching to ChPT of all (sub)amplitudes
 3. improved estimate of uncertainties
- Having evaluated all the following diagram




- the results for $\sigma(e^{+}e^{-} \rightarrow \pi^{+}\pi^{-}(\gamma))$ look as follows: **preliminary** J. Monnard thesis (2021)




Evaluation of $F_{\pi}^{V,\alpha}$

- Work in progress:
 1. $M_{\pi}^{+} - M_{\pi}^0$ effects missing
 2. controlled matching to ChPT of all (sub)amplitudes
 3. improved estimate of uncertainties
- Having evaluated all the following diagram



- the results for $\sigma(e^{+}e^{-} \rightarrow \pi^{+}\pi^{-}(\gamma))$ look as follows: **preliminary** J. Monnard thesis (2021)



Red curve corresponds to Hofer, Gluza, Jegerlehner (02)

- Ideally one would use the calculated RC directly in the data analysis

- to get an idea of the impact we did the following:

[thanks to M. Hoferichter and P. Stoffer]

1. remove RC from the measured $\sigma(e^+e^- \rightarrow \pi^+\pi^-(\gamma))$
2. fit with the dispersive representation for F_π^V
3. insert back the RC

- the impact on a_μ^{HVP} (comparison with result obtained by removing RC)

$$10^{11} \Delta a_\mu^{HVP} = \begin{cases} 10.2 \pm 0.5 \pm 5 & \text{sQED} \\ 10.5 \pm 0.5 & \text{triangle} \\ 13.2 \pm 0.5 & \text{full} \end{cases}$$

Preliminary, J. Monnard thesis (2021)

Introduction

Interference: RC to the forward-backward asymmetry in $e^+e^- \rightarrow \pi^+\pi^-$

Isospin-breaking corrections for $\pi\pi$ scattering

Dispersive approach to FSR in $e^+e^- \rightarrow \pi^+\pi^-$

Summary / Outlook

- **Dispersive** (pion-pole) determination of the **interference** terms to $e^+e^- \rightarrow \pi^+\pi^-$ and its contribution to the **forward-backward asymmetry** [Colangelo, Hoferichter, Monnard, JRE (2022)]
- Formalism for evaluating **dispersively RC** to the $\pi\pi$ scattering and F_π^V considering only **2π intermediate states** [Colangelo, Monnard, JRE (in progress)]
- our **preliminary** evaluation of the corrections to F_π^V shows **no** unexpectedly **large effects** [J. Monnard, PhD thesis, (2021)]
- our **preliminary** estimate of the impact on a_μ^{HVP} also shows **moderate effects** [J. Monnard, PhD thesis, (2021)]
- the final goal is to provide a **ready-to-use code** which can be implemented in MC and used in data analysis

Spare slides

- One-loop ChPT calculation

[Kaiser (2010)]

- Experimental results

[COMPASS (2012)]

- Dispersive result for the pion pole + resonances

[Colangelo, Monnard, JRE (in progress)]

