



FLOW-ORIENTED PERTURBATION THEORY

Collaborators: M. Borinsky, Z. Capatti and E. Laenen.

RADCOR 2023, Tuesday 30th May, 2023

Alexandre Salas-Bernárdez



Introduction I

low-oriented perturbation theory

The *p-x* representation of the S-m 0000000000 Unitarity and cut integrals

Outline

1 Introduction and motivation.

ow-oriented perturbation theory

The *p-x* representation of the S-m 0000000000 Unitarity and cut integrals

Outline

1 Introduction and motivation.

2 Derivation of FOPT.

ow-oriented perturbation theory

The *p-x* representation of the S-m 0000000000 Unitarity and cut integrals

Outline

1 Introduction and motivation.

2 Derivation of FOPT. Examples.

Outline

1 Introduction and motivation.

- 2 Derivation of FOPT. Examples.
- 3 Hybrid S-Matrix representation and the Flow polytope.

Outline

1 Introduction and motivation.

- 2 Derivation of FOPT. Examples.
- 3 Hybrid S-Matrix representation and the Flow polytope.
- 4 Unitarity and cut integrals in FOPT.

Based on "Flow-oriented perturbation theory", JHEP 01 (2023), 172 https://arxiv.org/abs/2210.05532.

Introduction

Alexandre Salas-Bernárdez

The *p-x* representation of the S-mat

Unitarity and cut integrals

3D representations of Feynman integrals



 Time Ordered Perturbation Theory (TOPT)



The *p-x* representation of the S-mat

Unitarity and cut integrals

3D representations of Feynman integrals

- Famous non-(manifestly)covariant approaches:
 - Time Ordered Perturbation Theory (TOPT)
 - Loop-tree duality.



Introduction Flow-oriented perturba

The *p*-*x* representation of the S-mat

Unitarity and cut integrals

Coordinate space formulation of QFTs

Coordinate space treatments:

- Unitarity and the Largest Time equation.
- Multi-loop renormalization group invariants.
- Factorization results.
- Axiomatic QFT.
- PDFs.
- · ...

The *p*-*x* representation of the S-mat

Unitarity and cut integrals

Coordinate space scalar QFT

$$\Delta_F(z) = \int \frac{d^4 p}{(2\pi)^4} e^{-ip \cdot z} \frac{i}{p^2 + i\epsilon} = \frac{1}{(2\pi)^2} \frac{1}{-z^2 + i\epsilon} \,.$$

The *p*-*x* representation of the S-mat

Unitarity and cut integrals

Coordinate space scalar QFT

$$\Delta_F(z) = \int \frac{d^4p}{(2\pi)^4} e^{-ip \cdot z} \frac{i}{p^2 + i\epsilon} = \frac{1}{(2\pi)^2} \frac{1}{-z^2 + i\epsilon} \,.$$

■ Scalar *n*-point Green's function

$$\begin{split} \mathsf{\Gamma}(x_1, ..., x_{|V^{\mathrm{ext}}|}) &= \left\langle 0 | \, \mathcal{T}(\varphi(x_1) \cdots \varphi(x_{|V^{\mathrm{ext}}|})) | 0 \right\rangle \\ &= \sum_{\mathcal{G}} \frac{1}{\mathsf{Sym} \, \mathcal{G}} \mathcal{A}_{\mathcal{G}}(x_1, \dots, x_{|V^{\mathrm{ext}}|}), \end{split}$$

The p-x representation of the S-ma 0000000000

Unitarity and cut integrals

Coordinate space scalar QFT

$$\Delta_F(z) = \int \frac{d^4p}{(2\pi)^4} e^{-ip \cdot z} \frac{i}{p^2 + i\epsilon} = \frac{1}{(2\pi)^2} \frac{1}{-z^2 + i\epsilon} \,.$$

■ Scalar *n*-point Green's function

$$\begin{split} \mathsf{\Gamma}(x_1, ..., x_{|V^{\mathrm{ext}}|}) &= \left\langle 0 | \, \mathcal{T}(\varphi(x_1) \cdots \varphi(x_{|V^{\mathrm{ext}}|})) | 0 \right\rangle \\ &= \sum_{G} \frac{1}{\mathsf{Sym} \; G} \mathcal{A}_{G}(x_1, \dots, x_{|V^{\mathrm{ext}}|}), \end{split}$$

• A graph G contributing to the Green's function

$$A_G(x_1,\ldots,x_{|V^{\text{ext}}|}) = \frac{(-ig)^{|V^{\text{int}}|}}{(2\pi)^{2|E|}} \left[\prod_{v \in V^{\text{int}}} \int d^4 y_v\right] \prod_{e \in E} \frac{1}{-z_e^2 + i\varepsilon}$$

The *p*-x representation of the S-ma

Unitarity and cut integrals

Coordinate space triangle diagram



$$\begin{split} \mathcal{A}_G(x_1, x_2, x_3) &= \frac{(-ig)^3}{(2\pi)^{12}} \int \left[\prod_{\nu \in V^{\text{int}}} \mathrm{d}^4 y_\nu \right] \times \\ &\times \frac{1}{(x_1 - y_1)^2 (x_2 - y_2)^2 (x_3 - y_3)^2 (y_1 - y_2)^2 (y_2 - y_3)^2 (y_1 - y_3)^2} \,, \end{split}$$

Flow-oriented perturbation theory

Alexandre Salas-Bernárdez

Performing time integrations

In the spirit of TOPT we perform $\left[\int dy_{\nu}^{0}\right]$ integrations to obtain a 3D representation of coordinate space diagrams:

- In doing so we introduce auxiliary energy variables.
- Perform Cauchy integrations.

The result is a sum over the different energy flows (orientations σ) in the diagram, with energies being conserved at each vertex.

$$\frac{\mathcal{A}_{\mathcal{G}}(x_1,\ldots,x_{|V^{\mathrm{ext}}|})}{\operatorname{\mathsf{Sym}} \mathcal{G}} = \sum_{\langle \boldsymbol{\sigma} \rangle} \frac{\mathcal{A}_{\mathcal{G},\boldsymbol{\sigma}}(x_1,\ldots,x_{|V^{\mathrm{ext}}|})}{\operatorname{\mathsf{Sym}}(\mathcal{G},\boldsymbol{\sigma})},$$

It is possible to resolve the energy integrations and conservation conditions for each orientation σ on a graph G in terms of "cycle" energy variables

It is possible to resolve the energy integrations and conservation conditions for each orientation σ on a graph G in terms of "cycle" energy variables







It is possible to resolve the energy integrations and conservation conditions for each orientation σ on a graph G in terms of "cycle" energy variables



Energy conservation in (G, σ) \Rightarrow Strongly connected closed graph (G°, σ)

It is possible to resolve the energy integrations and conservation conditions for each orientation σ on a graph G in terms of "cycle" energy variables:

$$egin{aligned} \mathcal{A}_{G, \pmb{\sigma}}(x_1, \dots, x_{|V^{ ext{ext}}|}) &\propto \left(\prod_{v \in V^{ ext{int}}} \int \mathrm{d}^3 ec{y}_v
ight) imes \ & imes \left(\prod_{e \in E} rac{1}{2|ec{z}_e|}
ight) \prod_{p \in ext{cycles}} rac{1}{\gamma_p + au_p + iarepsilon}. \end{aligned}$$

 τ_p is the time difference and γ_p the sum of the lengths of the edges passed in the cycle.

The *p*-*x* representation of the S-mat

Unitarity and cut integrals

One loop self energy graph



$$A_G(x_1, x_2) = \frac{(-ig)^2}{(4\pi^2)^4} \int d^4 y_1 d^4 y_2 \frac{1}{-z_1^2 + i\varepsilon} \frac{1}{-z_2^2 + i\varepsilon} \frac{1}{-z_3^2 + i\varepsilon} \frac{1}{-z_4^2 + i\varepsilon}$$

Flow-oriented perturbation theory The *p*-*x* representation of the S-matrix Unitarity and cut integrals

One loop self energy closed graph



Flow-oriented perturbation theory The *p*-*x* representation of the S-matrix Unitarity and cut integrals

One loop self energy closed graph



Next, draw all possible energy flows.

Introduction | 000000

Flow-oriented perturbation theory

The *p*-x representation of the S-mat

Unitarity and cut integrals

Energy flows through the closed bubble



Introduction Flo

Flow-oriented perturbation theory

The *p-x* representation of the S-mate

Unitarity and cut integrals

Energy flows through the closed bubble



Energy must be conserved at each vertex.

Introduction Flow

Flow-oriented perturbation theory

The *p*-*x* representation of the S-mat

Unitarity and cut integrals

Energy flows (= orientations) through the closed bubble



Flow-oriented perturbation theory The p-x representation of the S-matrix Unitarity and cut integrals

Energy flows (= orientations) through the closed bubble



Flow-oriented perturbation theory The *p*-*x* representation of the S-matrix Unitarity and cut integrals

3D representation of the bubble



The *p-x* representation of the S-mat

Unitarity and cut integrals

3D representation of the bubble



By collecting the overall and individual symmetry factors, we have that

$$\frac{1}{2}A(x_1, x_2) = \frac{1}{2}A_{G, \sigma_{(1)}} + A_{G, \sigma_{(2)}} + A_{G, \sigma_{(8)}} + \frac{1}{2}A_{G, \sigma_{(9)}}.$$

Flow-oriented perturbation theory The *p*-*x* representation of the S-matrix Unitarity and cut integrals



Flow-oriented perturbation theory The p-x representation of the S-matrix Unitarity and cut integrals





Flow-oriented perturbation theory The p-x representation of the S-matrix Unitarity and cut integrals





Flow-oriented perturbation theory The *p*-*x* representation of the S-matrix Unitarity and cut integrals



Introduction

Flow-oriented perturbation theory

The *p*-*x* representation of the S-mat

Unitarity and cut integrals



000000000000000

Flow-oriented perturbation theory The p-x representation of the S-matrix Unitarity and cut integrals

Decomposition of an orientation into cycles



UV divergent if y1->y2

Flow-oriented perturbation theory The *p*-*x* representation of the S-matrix Unitarity and cut integrals

 x_5

 $- x_4$

 x_3

Routes and cycles: UV singularities in FOPT

Two types of paths:


x_5

 $- x_4$

 x_3

Routes and cycles: UV singularities in FOPT

Two types of paths:



Routes and cycles: UV singularities in FOPT

Two types of paths:



UV singularities of cycles match those of the covariant Feynman diagrams.

 \Rightarrow Amplitudes can be regularised as "usual"

Routes and cycles: UV singularities in FOPT

Two types of paths:



UV singularities of cycles match those of the covariant Feynman diagrams.

 \Rightarrow Amplitudes can be regularised as "usual" (it is coordinate space).

Flow-oriented perturbation theory The p-x representation of the S-matrix Unitarity and cut integrals

Long and finite distance singularities in FOPT

Diagrams in FOPT fail to reproduce the finite distance (collinear) and long distance (soft) divergent behaviour expected from momentum space results.

Long and finite distance singularities in FOPT

Diagrams in FOPT fail to reproduce the finite distance (collinear) and long distance (soft) divergent behaviour expected from momentum space results.

 \Rightarrow We shift our attention to the S-matrix and construct an FOPT representation of it.

The *p*-*x* representation of the S-matrix

Alexandre Salas-Bernárdez

Hybrid representation of the S-matrix

Hybrid representation of the S-matrix

$$S(\{p_i\}_{i \in V_{\text{in}}^{\text{ext}}}, \{p_f\}_{f \in V_{\text{out}}^{\text{ext}}}) = Z^{|V_{\text{ext}}|/2} \widetilde{\Gamma}_T(\{p_a\}_{a \in V^{\text{ext}}}),$$

Hybrid representation of the S-matrix

$$S(\{p_i\}_{i\in V_{\mathrm{in}}^{\mathrm{ext}}}, \{p_f\}_{f\in V_{\mathrm{out}}^{\mathrm{ext}}}) = Z^{|V_{\mathrm{ext}}|/2}\widetilde{\Gamma}_{\mathcal{T}}(\{p_a\}_{a\in V^{\mathrm{ext}}}),$$

$$\widetilde{\Gamma}_{\mathcal{T}}(\{p_{a}\}_{a\in V^{\mathrm{ext}}}) = \left[\prod_{i\in V_{\mathrm{in}}^{\mathrm{ext}}}\widetilde{\Delta}_{R}(p_{i})\right]^{-1} \left[\prod_{f\in V_{\mathrm{out}}^{\mathrm{ext}}}\widetilde{\Delta}_{A}(p_{f})\right]^{-1} \widetilde{\Gamma}(\{p_{a}\}_{a\in V^{\mathrm{ext}}}).$$

Hybrid representation of the S-matrix

$$S(\{p_i\}_{i\in V_{\mathrm{in}}^{\mathrm{ext}}}, \{p_f\}_{f\in V_{\mathrm{out}}^{\mathrm{ext}}}) = Z^{|V_{\mathrm{ext}}|/2}\widetilde{\Gamma}_{\mathcal{T}}(\{p_a\}_{a\in V^{\mathrm{ext}}}),$$

$$\widetilde{\Gamma}_{\mathcal{T}}(\{p_{a}\}_{a\in V^{\text{ext}}}) = \left[\prod_{i\in V_{\text{in}}^{\text{ext}}}\widetilde{\Delta}_{R}(p_{i})\right]^{-1} \left[\prod_{f\in V_{\text{out}}^{\text{ext}}}\widetilde{\Delta}_{A}(p_{f})\right]^{-1} \widetilde{\Gamma}(\{p_{a}\}_{a\in V^{\text{ext}}}).$$
$$\widetilde{\Gamma}(\{p_{a}\}_{a\in V^{\text{ext}}}) = \int \left[\prod_{a\in V^{\text{ext}}} d^{4}x_{a} e^{ix_{a} \cdot p_{a}}\right] \Gamma(\{x_{a}\}_{a\in V^{\text{ext}}}),$$

Introduction Flow-oriented perturbation theory October October

Hybrid representation of the S-matrix

Next we use FOPT representation of the Green's function

$$\widetilde{\Gamma}(\{p_a\}_{a\in V^{\mathrm{ext}}}) = \int \left[\prod_{a\in V^{\mathrm{ext}}} \mathrm{d}^4 x_a \, e^{ix_a \cdot p_a}\right] \Gamma(\{x_a\}_{a\in V^{\mathrm{ext}}}),$$

Hybrid representation of the S-matrix

Next we use FOPT representation of the Green's function

$$\widetilde{\Gamma}(\{p_a\}_{a\in V^{\mathrm{ext}}}) = \int \left[\prod_{a\in V^{\mathrm{ext}}} \mathrm{d}^4 x_a \, e^{i x_a \cdot p_a}\right] \Gamma(\{x_a\}_{a\in V^{\mathrm{ext}}}),$$

$$\widetilde{\Gamma}(\{p_a\}_{a\in V^{\text{ext}}}) = \sum_{(G,\sigma)} \frac{1}{\text{Sym}(G,\sigma)} \widetilde{A}_{G,\sigma}(\{p_a\}_{a\in V^{\text{ext}}}),$$

Hybrid representation of the S-matrix

Next we use FOPT representation of the Green's function

$$\widetilde{\Gamma}(\{p_a\}_{a\in V^{\mathrm{ext}}}) = \int \left[\prod_{a\in V^{\mathrm{ext}}} \mathrm{d}^4 x_a \, e^{ix_a \cdot p_a}\right] \Gamma(\{x_a\}_{a\in V^{\mathrm{ext}}}),$$

$$\widetilde{\Gamma}(\{p_a\}_{a\in V^{\text{ext}}}) = \sum_{(G,\sigma)} \frac{1}{\text{Sym}(G,\sigma)} \widetilde{A}_{G,\sigma}(\{p_a\}_{a\in V^{\text{ext}}}),$$

where the Fourier transform of a FOPT orientation is given by

$$\widetilde{A}_{G,\sigma}(\{p_a\}_{a\in V^{\text{ext}}}) = \int \left[\prod_{a\in V^{\text{ext}}} \mathrm{d}^4 x_a \, e^{ix_a \cdot p_a}\right] A_{G,\sigma}(\{x_a\}_{a\in V^{\text{ext}}}).$$

p-x representation of the S-matrix

It is possible to perform the Fourier transform explicitly and the final result equals

$$S(\{p_i\}_{i\in V_{\text{in}}^{\text{ext}}}, \{p_f\}_{f\in V_{\text{out}}^{\text{ext}}}) = \sum_{(G,\sigma)} \frac{S_{G,\sigma}(\{p_i\}_{i\in V_{\text{in}}^{\text{ext}}}, \{p_f\}_{f\in V_{\text{out}}^{\text{ext}}})}{\text{Sym}(G,\sigma)},$$

p-x representation of the S-matrix

It is possible to perform the Fourier transform explicitly and the final result equals

$$S(\{p_i\}_{i\in V_{\text{in}}^{\text{ext}}}, \{p_f\}_{f\in V_{\text{out}}^{\text{ext}}}) = \sum_{(G,\sigma)} \frac{S_{G,\sigma}(\{p_i\}_{i\in V_{\text{in}}^{\text{ext}}}, \{p_f\}_{f\in V_{\text{out}}^{\text{ext}}})}{\text{Sym}(G,\sigma)},$$

$$\begin{split} S_{G,\sigma} &\propto \delta \left(\sum_{\boldsymbol{a} \in V^{\text{ext}}} p_{\boldsymbol{a}}^{0} \right) \times \\ &\times \int \frac{\left[\prod_{\boldsymbol{v} \in V^{\text{int}}} \mathrm{d}^{3} \vec{y}_{\boldsymbol{v}} \right] \left[\prod_{\boldsymbol{a} \in V^{\text{ext}}} e^{-i \vec{y}_{\overline{a}} \cdot \vec{p}_{\boldsymbol{a}}} \right]}{\left[\prod_{\boldsymbol{e} \in F^{\text{int}}} 2 |\vec{z}_{\boldsymbol{e}}| \right] \left[\prod_{c \in \Gamma^{\text{int}}} \gamma_{c} \right]} \widehat{\mathcal{F}}_{G,\sigma}^{\{p_{\boldsymbol{a}}^{0}\}}(\boldsymbol{\gamma}^{t} + i\varepsilon 1). \end{split}$$

p-x representation of the S-matrix

It is possible to perform the Fourier transform explicitly and the final result equals

$$S(\{p_i\}_{i\in V_{\text{in}}^{\text{ext}}}, \{p_f\}_{f\in V_{\text{out}}^{\text{ext}}}) = \sum_{(G,\sigma)} \frac{S_{G,\sigma}(\{p_i\}_{i\in V_{\text{in}}^{\text{ext}}}, \{p_f\}_{f\in V_{\text{out}}^{\text{ext}}})}{\text{Sym}(G,\sigma)},$$

$$\begin{split} S_{G,\sigma} &\propto \delta \left(\sum_{a \in V^{\mathrm{ext}}} p_a^0 \right) \times \\ & \times \int \frac{\left[\prod_{v \in V^{\mathrm{int}}} \mathrm{d}^3 \vec{y}_v \right] \left[\prod_{a \in V^{\mathrm{ext}}} e^{-i\vec{y}_{\overline{a}} \cdot \vec{p}_a} \right]}{\left[\prod_{e \in E^{\mathrm{int}}} 2|\vec{z}_e| \right] \left[\prod_{c \in \Gamma^{\mathrm{int}}} \gamma_c \right]} \widehat{\mathcal{F}}_{G,\sigma}^{\{p_a^0\}}(\gamma^t + i\varepsilon 1). \end{split}$$



The flow polytope

$$\widehat{\mathcal{F}}_{G,\sigma}^{\{\boldsymbol{p}_a^0\}}(\boldsymbol{\gamma}+i\varepsilon 1) = \int_{\mathcal{F}_{G,\sigma}^{\{\boldsymbol{p}_a^0\}}} \mathrm{d}\boldsymbol{E} \ e^{i\boldsymbol{E}\cdot(\boldsymbol{\gamma}+i\varepsilon 1)}.$$

The flow polytope

$$\widehat{\mathcal{F}}_{G,\sigma}^{\{p_a^0\}}(\gamma+i\varepsilon 1) = \int_{\mathcal{F}_{G,\sigma}^{\{p_a^0\}}} \mathrm{d}\boldsymbol{E} \ e^{i\boldsymbol{E}\cdot(\gamma+i\varepsilon 1)}.$$

 $\mathcal{F}_{G,\sigma}^{\{p_a^o\}}$ is swept out by all tuples $(E_r)_{r\in\Gamma^{ext}}$ which fulfill

$$E_{
m r} \ge 0$$
 for all $m r \in \Gamma^{
m ext}$,
 $\sum_{
m r
i i} E_{
m r} = -p_i^0$ for all $i \in V_{
m in}^{
m ext}$,
 $\sum_{
m r
i f} E_{
m r} = -p_f^0$ for all $f \in V_{
m out}^{
m ext}$.

The flow polytope

$$\widehat{\mathcal{F}}_{G,\sigma}^{\{p_a^0\}}(\boldsymbol{\gamma}+i\varepsilon 1) = \int_{\mathcal{F}_{G,\sigma}^{\{p_a^0\}}} \mathrm{d}\boldsymbol{E} \, e^{i\boldsymbol{E}\cdot(\boldsymbol{\gamma}+i\varepsilon 1)}.$$

 $\mathcal{F}_{G\sigma}^{\{p_{a}^{a}\}}$ is swept out by all tuples $(E_{r})_{r\in\Gamma^{\mathrm{ext}}}$ which fulfill

$$E_{
m r} \ge 0$$
 for all $m r \in \Gamma^{
m ext}$,
 $\sum_{
m r \ni i} E_{
m r} = -p_i^0$ for all $i \in V_{
m in}^{
m ext}$,
 $\sum_{
m r \ni f} E_{
m r} = -p_f^0$ for all $f \in V_{
m out}^{
m ext}$.

Nice features regarding the cancellation of spurious singularities.

Example: The p-x representation of the triangle diagram



Example: The *p*-*x* representation of the triangle diagram





 $E_1+E_3=-p_2^0\;$ The flow polytope is cut out by the conditions:

$$egin{aligned} & E_1, E_2, E_3 \geq 0\,, \ & E_1 + E_2 + E_3 = p_1^0\,, \ & E_1 + E_3 = -p_2^0\,, \ & E_2 = -p_3^0\,, \end{aligned}$$

I

IR singularities in the p-x representation

Collinear singularities are studied taking limits as $(\lambda \rightarrow \infty)$:



Unitarity and cut integrals

IR singularities in the p-x representation

Collinear singularities are studied taking limits as $(\lambda \to \infty)$:



IR singularities in the p-x representation

Collinear singularities are studied taking limits as $(\lambda \to \infty)$:



We find a per-diagram factorization of collinear and hard singularities!

Per-diagram factorization



only two-cut yield collinear singularities

Per-diagram factorization of the S-matrix IR singularities

$$\begin{split} s_{G,\sigma}(\{p_1,\ldots,p_k\},\{p_{k+1},\ldots,p_n\}) &= \\ &= -\frac{2\pi i}{4}\log\frac{p_n^2}{Q^2}\int_0^1 \mathrm{d} x\,s_{(G,\sigma)_{\mathsf{hard}}}s_{(G,\sigma)_{\mathsf{col}}} + \mathcal{O}_{p_n^2 \to 0}(1)\,, \end{split}$$



Soft-collinear singularity in the triangle diagram

We can study in the triangle the overlap of the collinear singularity with the soft singularity



Soft-collinear singularity in the triangle diagram

We can study in the triangle the overlap of the collinear singularity with the soft singularity



Soft-collinear singularity in the triangle diagram

We can study in the triangle the overlap of the collinear singularity with the soft singularity



 \Rightarrow appearance of double-log Sudakov logs:

$$s_{G,\sigma}(\{p_1\},\{p_2,p_3\}) = -i\frac{(2\pi)^2}{8}\frac{\log\frac{p_2^2}{p_1^2}\log\frac{p_3^2}{p_1^2}}{p_2\cdot p_3} + \mathcal{O}_{\substack{p_2^2\to 0\\p_3^2\to 0}}(1).$$

Unitarity and cut integrals

Alexandre Salas-Bernárdez

Introduction Flow-oriented perturbation theory

The *p-x* representation of the S-mat

Unitarity and cut integrals $_{\bigcirc \odot \odot \odot}$

Cuts relating virtual and real processes



Introduction Flow-oriented perturbation theory

The *p-x* representation of the S-mat

Unitarity and cut integrals

Cuts relating virtual and real processes

$$2 \operatorname{Im} \left[\begin{array}{c} a \\ \end{array} \right] = \sum_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}[\begin{array}[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}[\begin{array}[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}[\begin{array}[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\begin{array}[\begin{array}[\begin{array}{c} a \\ \end{array} \right] = \int_{r} \int d\Pi_{f} \left[\left[\begin{array}[\begin{array}[\begin{array}{c} a \end{array} \right] = \int_{r}$$

Loop Tree Duality puts all virtual and real corrections to a cross section under the same integral sign.

0000

FOPT cut integrals

It is possible to extend FOPT to cut integrals.



A remarkable property arises: different sized cuts have the same integral measure.

00.00

FOPT cut integrals

It is possible to extend FOPT to cut integrals.



A remarkable property arises: different sized cuts have the same integral measure.

 \Rightarrow Advantage: IR singularities in numerical evaluations will cancel locally (no need for Loop Tree Duality).

Outlook

■ FOPT offers promising features:

Canonical Feynman rules.

Outlook

■ FOPT offers promising features:

- Canonical Feynman rules.
- 3D representation of the S-matrix and the flow polytope.
Outlook

FOPT offers promising features:

- Canonical Feynman rules.
- 3D representation of the S-matrix and the flow polytope.
- Per-diagram factorization of IR singularities in the S-matrix.

Next steps are:

,

Extend FOPT to D dimensions.

Outlook

FOPT offers promising features:

- Canonical Feynman rules.
- 3D representation of the S-matrix and the flow polytope.
- Per-diagram factorization of IR singularities in the S-matrix.

Next steps are:

,

- Extend FOPT to D dimensions.
- Extend it to massive and fermion lines.

```
Introduction Flow-oriented perturbation theory The p-x representation of the S-matrix Unitarity and cut integrals
```

Outlook

FOPT offers promising features:

- Canonical Feynman rules.
- 3D representation of the S-matrix and the flow polytope.
- Per-diagram factorization of IR singularities in the S-matrix.

Next steps are:

,

- Extend FOPT to D dimensions.
- Extend it to massive and fermion lines.
- Factorization.