Renormalization of twist-two operators and new 4-loop results for  $N_f^2$  pure-singlet splitting functions

Tong-Zhi Yang

with Thomas Gehrmann, Andreas von Manteuffel and Vasily Sotnikov Based on 2302.00022 [JHEP 04 (2023) 041] and a forthcoming paper

RADCOR 2023, Crieff Hydro Hotel, 30 May 2023









Tong-Zhi Yang (University of Zurich) Renormalization of twist-two operators

### Parton densities and splitting functions





$$x_B = \frac{-q^2}{2P \cdot q}$$

Factorization

$$\sigma \sim \sum_{a} f_{a|N}(x_B) \otimes \hat{\sigma}_a(x_B)$$

• Quark parton density in axial gauge

$$f_{q|N}(x_B) = \int rac{dt}{2\pi} e^{-i x_B t \Delta \cdot P} \langle N(P) | ar{\psi}(t\Delta) rac{A}{2} \psi(0) | N(P) 
angle, \ \Delta^2 = 0$$

• Splitting functions (SFs) govern the DGLAP evolutions of PDFs

$$\frac{df_{i|N}}{d\ln\mu} = 2\sum_{k} \frac{P_{ik}}{k} \otimes f_{k|N}$$

#### Why 4-loop SFs for the evolutions of N3LO PDFs?

• Expand PDFs and SFs with  $a_s = \alpha_s/(4\pi)$ 

$$f_{i|N} = f_{i|N}^{(0)} + f_{i|N}^{(1)} a_s + \dots + f_{i|N}^{(3)} a_s^3 + \dots$$
$$P_{ij} = P_{ij}^{(0)} a_s + \dots + P_{ij}^{(3)} a_s^4 + \dots$$

Evolution of *a<sub>s</sub>*

$$\frac{da_s}{d\ln\mu} = -2(a_s^2\beta_0 + a_s^3\beta_1 + \cdots)$$

A consistent evolution of N3LO PDFs requires 4-loop SFs

$$f_{i|N}^{(3)} \frac{da_s^3}{d\ln\mu} = f_{i|N}^{(3)} (-6a_s^4\beta_0 + \cdots) = \sum_k \frac{P_{ik}^{(3)}}{a_s^4} a_s^4 \otimes f_{k|N}^{(0)} + \cdots$$

#### Motivations for four-loop splitting functions

- Several  $\hat{\sigma}$  are available at N3LO, but N3LO PDFs are missing
- The fields in fitting N3LO PDFs are active
  - MSHT20 aN3LO talks by Thomas Cridge, Lucian Harland-Lang from DIS2023
  - NNPDF in progress towards aN3LO talk by Giacomo Magni from DIS2023
  - CT are planning talk by Pavel Nadolsky from DIS2023



- Scale uncertainty at N3LO using NNLO PDF remains at 1% level[Chen,Gehrmann,Glover,Husss,Yang,Zhu,2021]
- Seems that aN3LO PDF introduces large corrections to  $\sigma$  talk by Tobias Neumman from DIS2023

Tong-Zhi Yang (University of Zurich) Renormalization of twist-two operators

#### Splitting functions & Anomalous dimensions

Mellin transformation

$$f_q(n) = -\int_0^1 dx_B \; x_B^{n-1} f_q(x_B) \, , \gamma_{ij}(n) = -\int_0^1 dx_B \; x_B^{n-1} P_{ij}(x_B)$$

• DGLAP evolution in *n*-space

$$\frac{d}{d\ln\mu}f_q(n,\mu^2) = -2\sum_j \gamma_{qj}(n) f_j(n,\mu^2)$$

• PDFs in *n*-space are hadronic operator matrix elements (OMEs)

$$f_q(n) \sim \langle N(P) | \bar{\psi} \Delta (\Delta \cdot D)^{n-1} \psi | N(P) \rangle$$

#### Twist-two operators

According to the flavor group,

• Non-singlet: a single operator

$$O_{q,k} = \frac{i^{n-1}}{2} \left[ \bar{\psi}_i \measuredangle (\Delta \cdot D)_{ij}^{n-1} \frac{\lambda_k}{2} \psi_j \right], k = 3, 8, \cdots n_f^2 - 1$$

 $\lambda_k/2$  is the diagonal generator of the flavor group

Singlet: two operators

$$O_q = \frac{i^{n-1}}{2} \left[ \bar{\psi}_i \measuredangle (\Delta \cdot D)_{ij}^{n-1} \psi_j \right],$$
  
$$O_g = -\frac{i^{n-2}}{2} \left[ \Delta_{\mu_1} G_{a,\mu}^{\mu_1} (\Delta \cdot D)_{ab}^{n-2} \Delta_{\mu_n} G_b^{\mu_n \mu} \right]$$

 $G_a^{\mu
u}$  is the gluon field strength tensor.

#### Renormalization of twist-two operators

• The non-singlet operator  $O_{q,k}$  is multiplicatively renormalized,

$$O_{q,k}^{\mathsf{R}} = Z^{\mathsf{ns}} O_{q,k}^{\mathsf{B}}$$

• The two singlet operators mix under renormalization,

$$\left(\begin{array}{c}O_{q}\\O_{g}\end{array}\right)^{\mathsf{R, naive}} = \left(\begin{array}{c}Z_{qq} & Z_{qg}\\Z_{gq} & Z_{gg}\end{array}\right) \left(\begin{array}{c}O_{q}\\O_{g}\end{array}\right)^{\mathsf{B}}$$

• Evolution equation for the renormalization constants

$$\frac{dZ_{ij}}{d\ln\mu} = -2\sum_{k=q,g} \frac{\gamma_{ik}(n)Z_{kj}}{\gamma_{ik}(n)Z_{kj}}$$

Extract anomalous dimensions from the renormalization constants

$$Z_{ij} = \delta_{ij} + \sum_{l=1}^{\infty} a_s^l \frac{1}{l \epsilon} \gamma_{ij}^{(k-1)} + \cdots$$

## DIS method vs OME method

• Forward DIS (gauge invariant)



- Off-shell OMEs are not gauge invariant (due to off-shell external gluons), physical operators mix with unknown gauge-variant (GV) operators
- Main goal: find all GV operators or their Feynman rules

Once all GV operators are known, the off-shell OME method can be used to determine SFs efficiently

#### A bit of history about the calculations of SFs

- The first one-loop results are from off-shell OMEs
  - ▶ Non-singlet and singlet [D.J. Gross, F. Wilczek, 1973, 1974]
- Two-loop results
  - Non-singlet from off-shell OMEs[E.G. Floratos et al. 1977]
  - Singlet: inconsistences from off-shell OMEs in covariant gauge (Flaws due to omitting GV operators)[E.G. Floratos et al. 1978] and from off-shell OMEs in axial gauge (Correct) [Furmanski and Petronzio, 1980]
- The first three-loop results are from DIS
  - Non-singlet and Singlet[Moch, Vermaseren and Vogt, 2004,2004]
- Partial four-loop results
  - ▶ Non-singlet with  $n \le 16$  from off-shell OMEs[S. Moch et al. 2017]
  - Singlet with  $n \leq 8$  from DIS[S. Moch et al. 2021]
  - ▶ Pure singlet with n ≤ 20 from off-shell OMEs[G. Falcioni et al. 2023] see the talk by S. Moch

#### Only a few low-n results at four-loop are available

#### Significant efforts in deriving GV operators

- [D.J. Gross, F. Wilczek, 1974] pointed out possible mixing with GV operators
- [J.A. Dixon and J.C. Taylor, 1974] constructed order  $g_s$  GV operators, not clear how to generalize to higher order
- [Joglekar and Lee, 1975] gave a general theorem about the renormalization of gauge invariant operators No explicit results were given
- [J. C. Collins and R. J. Scalise, 1994] studied the renormalization of energy-momentum tensor in detail and pointed out subtleties of theorem by Joglekar and Lee
- [G. Falcioni and F. Herzog, 2022] constructed the GV operators for a fixed *n* based on a generalized BRST symmetry. Promising, however more and more number of operators are needed for higher *n*.
- This talk: A new framework which enables the derivation of all-*n* GV operator (counterterm) Feynman rules to any loop orders

#### A new framework of deriving GV operators

- Guiding principles:
  - A twist-two operator has infinite mass dimension when  $n 
    ightarrow \infty$
  - Infinite GV operators are required to renormalize the physical operators
  - Some GV operators only contribute starting at higher-loop order
- Extend the naive renormalization of the operator Og,

$$O_{g}^{\mathsf{R}} = Z_{gq}O_{q}^{\mathsf{B}} + Z_{gg}O_{g}^{\mathsf{B}} + Z_{gA}\left(O_{A}^{\mathsf{B}} + O_{B}^{\mathsf{B}} + O_{C}^{\mathsf{B}}\right) + [ZO]_{g}^{\mathsf{GV}}$$
$$Z_{gA} = \mathcal{O}(a_{s}), \ [ZO]_{g}^{\mathsf{GV}} = \sum_{l=2}^{\infty} a_{s}^{l} \ [ZO]_{g}^{\mathsf{GV}, \ (l)}$$

•  $O_A$  (gluon fields only),  $O_B$ (quark+gluon fileds),  $O_C$  (ghost + gluon fields).  $[ZO]_g^{\text{GV}, (l)}$ : collection of counterterms

#### Generalized mixing matrix

• The framework can be applied to the renormalization of  $O_q$  directly

$$O_q^{\mathsf{R}} = Z_{qq}O_q^{\mathsf{B}} + Z_{qg}O_g^{\mathsf{B}} + Z_{qA}\left(O_A^{\mathsf{B}} + O_B^{\mathsf{B}} + O_C^{\mathsf{B}}\right) + [ZO]_q^{\mathsf{GV}}$$

where  $Z_{qA} = \mathcal{O}(\alpha_s^2), [ZO]_q^{\mathsf{GV}} = \mathcal{O}(\alpha_s^3)$ 

Generalized mixing matrix

$$\left(\begin{array}{c}O_{q}\\O_{g}\\O_{ABC}\end{array}\right)^{\mathsf{R}} = \left(\begin{array}{cc}Z_{qq} & Z_{qg} & Z_{qA}\\Z_{gq} & Z_{gg} & Z_{gA}\\0 & 0 & Z_{AA}\end{array}\right) \left(\begin{array}{c}O_{q}\\O_{g}\\O_{ABC}\end{array}\right)^{\mathsf{B}} + \left(\begin{array}{c}[ZO]_{q}^{\mathrm{GV}}\\[ZO]_{g}^{\mathrm{GV}}\\[ZO]_{A}\end{bmatrix}\right)^{\mathsf{B}}$$

• Where  $O_{ABC} = O_A + O_B + O_C$ 

 The renormalizations of GV operators don't mix with physical operators, compatible with the theorems given by[Joglekar and Lee, 1975]

D

#### Derive Feynman rules from off-shell OMEs

- Idea: derive Feynman rules instead of GV operators themselves
- Consider all off-shell OMEs with 2j + m-gluon external states

$$\begin{split} \langle j|O_g^{\mathsf{R}}|j+mg\rangle_{1\mathsf{Pl}}^{\mu_1\cdots\mu_m} &= \langle j|(Z_{gq}O_q^{\mathsf{B}}+Z_{gg}O_g^{\mathsf{B}})|j+mg\rangle_{1\mathsf{Pl}}^{\mu_1\cdots\mu_m} \\ &+ \langle j|Z_{gA}O_{ABC}^{\mathsf{B}}|j+mg\rangle_{1\mathsf{Pl}}^{\mu_1\cdots\mu_m} + \langle j|\left[ZO\right]_g^{\mathsf{GV}}|j+mg\rangle_{1\mathsf{Pl}}^{\mu_1\cdots\mu_m}, \, j=q,g \text{ or } \mathsf{c} \end{split}$$

• Expand OMEs order by order in loops and legs

$$\langle j|O|j+mg\rangle^{\mu_1\cdots\mu_m} = \sum_{l=1}^{\infty} \left[ \langle j|O|j+mg\rangle^{\mu_1\cdots\mu_m,\,(l),\,(m)} \right] \left(\frac{\alpha_s}{4\pi}\right)^l g_s^m$$

- Left: UV renormalized and IR finite  $\rightarrow$  no poles in  $\epsilon$
- Right: Each term is UV divergent, but the sum should be finite
  - Requirement of the finiteness allows to determine couterterm Feynman rules of unknown GV operators order by order

#### Determine Feynman rules for O<sub>ABC</sub>

• As an example, consider all off-shell two ghosts + *m*-gluon external states and expand to one-loop order

$$Z_{g\!A}^{(1)} \left\langle c | O_C | c + m \, g \right\rangle_{1\mathsf{Pl}}^{\mu_1 \cdots \mu_m, \, (0), \, (m)} = - \left[ \left\langle c | O_g | c + m \, g \right\rangle_{1\mathsf{Pl}}^{\mu_1 \cdots \mu_m, \, (1), \, (m), \, \mathsf{B}} \right]_{1/\epsilon}$$

•  $Z_{gA}^{(1)}$  is a *m*-independent constant and can be determined from m=0

$$Z_{gA}^{(1)} = \frac{-C_A}{\epsilon} \frac{1}{n(n-1)}$$



Sample digram to extract Feynman rules for  $O_C$  with m=2

# Determine Feynman rules for $[ZO]_g^{\text{GV},(2)}$

• As an example, consider all off-shell two ghosts + *m*-gluon external states and expand to two-loop order

$$\begin{split} &\langle c|\left[ZO\right]_{g}^{\mathrm{GV},\,(2)}\left|c+mg\right\rangle_{1\mathrm{Pl}}^{\mu_{1}\cdots\mu_{m},\,(0),\,(m)}=-\left\{\left[\langle c|O_{g}|c+mg\rangle_{1\mathrm{Pl}}^{\mu_{1}\cdots\mu_{m},\,(2),\,(m),\,\mathrm{B}}\right.\right.\\ &\left.+\left(Z_{c}^{(1)}+\frac{mZ_{g}^{(1)}}{2}+Z_{gg}^{(1)}-\frac{\beta_{0}(m+2)}{2\epsilon}\right)\langle c|O_{g}|c+mg\rangle_{1\mathrm{Pl}}^{\mu_{1}\cdots\mu_{m},\,(1),\,(m),\,\mathrm{B}}\right.\\ &\left.+Z_{gA}^{(1)}\left\langle c|O_{AC}|c+mg\rangle_{1\mathrm{Pl}}^{\mu_{1}\cdots\mu_{m},\,(1),\,(m),\,\mathrm{B}}+\cdots\right]_{\mathrm{div}}\right\} \end{split}$$



Sample digrams to extract Feynman rules for  $[ZO]_g^{
m GV,\,(2)}$  with m=1

# Three-loop singlet splitting functions from off-shell OMEs

### Sample Feynman diagrams

• Two-point diagrams with physical operators insertion



• Multi-point diagrams to infer GV counterterm Feynman rules





• Two-point diagrams with GV counterterm insertions



#### Computational methods

- Non-standard terms appearing in the Feynman rules
- Example: Feynman rules for  $O_q$  at lowest order

$$\xrightarrow{p_1,i_1} \xrightarrow{p_2,i_2} \rightarrow \not\Delta \left( \Delta \cdot p_1 \right)^{n-1}$$

 Sum the non-standard term into a linear propagator using a tracing parameter x, first proposed in[J. Ablinger, J. Blumlein, A. Hasselhuhn, S. Klein, C. Schneider, and F. Wissbrock, 2012] See also the talk by J. Bluemlein

$$(\Delta \cdot p)^{n-1} \to \sum_{n=1}^{\infty} x^n (\Delta \cdot p)^{n-1} = \frac{x}{1 - x\Delta \cdot p}$$

• Work in *x*-space throughout, take the coefficient of *x<sup>n</sup>* in the end using the package HarmonicSums[Ablinger 2010–].

#### Computational procedure

- Generate all relevant Feynman diagrams: QGRAF
- Substitute the *x*-space Feynman rules: Mathematica
- Evaluate dirac matrix and color algebra: FORM and Color
- Classify the topologies: self-written code, Reduze 2, FeynCalc
- Perform IBP reductions: FIRE6 with LiteRed, Reduze 2, Kira
- Derive differential equations with respect to the parameter x
- Solve the DEs
  - Two-point OMEs: in terms of special functions: the DEs are turned into canonical form proposed by Henn using CANONICA and Libra
  - Multi-point OMEs: expand the DEs to high orders in  $x \rightarrow 0$  limit and then reconstruct the final results based on heuristic ansatzs

#### Sample results: Feynman rules for $O_B$



Feynman rules for  $O_B$  operator with all momenta flowing into the vertex

#### Sample results for two-loop counterterm Feynman rules

 $\underbrace{p_{1}, \mu_{1}, a_{1}}_{p_{3}, \mu_{3}, a_{3}} \underbrace{(q, 2)}_{p_{2}, \mu_{2}, a_{2}}$ 

$$\begin{split} & \to 2ig_{s}C_{A}^{2}f^{a_{1}a_{2}a_{3}}\frac{1+(-1)^{n}}{256n(n-1)}\frac{\left(\Delta\cdot p_{1}\right)^{n-2}}{\Delta\cdot p_{2}}\bigg(-\Delta^{\mu_{1}}\Delta^{\mu_{2}}\Delta^{\mu_{3}}\sum_{i=1}^{3}p_{i}^{2} \\ & +\Delta^{\mu_{2}}\Delta^{\mu_{3}}p_{1}^{\mu_{1}}\Delta\cdot p_{1}+\Delta^{\mu_{1}}\Delta^{\mu_{3}}p_{2}^{\mu_{2}}\Delta\cdot p_{2}+\Delta^{\mu_{1}}\Delta^{\mu_{2}}p_{3}^{\mu_{3}}\Delta\cdot p_{3}\bigg) \\ & \times\bigg\{\frac{F_{-2,0}+(1-\xi)F_{-2,1}}{\epsilon^{2}}+\frac{F_{-1,0}+(1-\xi)F_{-1,1}}{\epsilon}\bigg\} \end{split}$$

•  $\epsilon$ -dependent and  $\xi$ -dependent

•  $F_{-1,0}$  contains generalized harmonic sums to weight-2, for example

$$S_{1,1}(1, z_1 + 1; n) = \sum_{t_1=1}^{n} \frac{1}{t_1} \sum_{t_2=1}^{t_1} \frac{(1+z_1)^{t_2}}{t_2}$$

• Weight-3 polylogarithms in *x*-space

#### Two-point OMEs with two-loop counterterm insertions

- For a fixed *n*, normal IBP, but need to reduce integrals with very high numerator degree
- All-n, IBP reduction with polylogarithms?
- Consider a general term of two-loop counterterms with 3-gluon vertex

$$ig_s f^{a_1 a_2 a_3} C_A^2 \Delta^{\mu_1} \Delta^{\mu_2} \Delta^{\mu_3} p_1^2 \sum_{m=0}^{n-3} a_{mn} (\Delta \cdot p_1)^m (\Delta \cdot p_2)^{n-3-m} + \cdots$$

where  $a_{mn}$  is known only for fixed m, n

• New idea: replace  $a_{mn}$  by another tracing parameter t

$$h(x,t) = \sum_{n=3}^{\infty} x^n \sum_{m=0}^{n-3} t^m (\Delta \cdot p_1)^m (\Delta \cdot p_2)^{n-3-m} = \frac{x^3}{(1 - x t \Delta \cdot p_1)(1 - x \Delta \cdot p_2)}$$

• Insert *h* into two-point diagrams:  $\langle g|h(x,t)|g\rangle = \sum_{n=3}^{\infty} x^n \sum_{m=0}^{n-3} t^m c_{mn}$ •  $\langle g|\sum_{m=0}^{n-3} a_{mn} (\Delta \cdot p_1)^m (\Delta \cdot p_2)^{n-3-m}|g\rangle = \sum_{m=0}^{n-3} a_{mn} c_{mn}$ 

Evaluate OMEs to any fixed n efficiently and reconstruct the full-n results

#### Results

- Combine results for all two-point OMEs (with  $O_q, O_g$  and GV counterterm insertions)
- Splitting functions: confirm the ξ-independence explicitly for the first time and recover the well known results in the literature
  - non-singlet:

$$\gamma^{(2)}_{\rm ns} - \gamma^{(2)}_{\rm ns} [{\rm MVV}] = 0$$

singlet:

$$\begin{split} \gamma_{qq}^{(2)} &- \gamma_{qq}^{(2)} [\mathsf{VMV}] = \mathbf{0} \,, \gamma_{qg}^{(2)} - \gamma_{qg}^{(2)} [\mathsf{VMV}] = \mathbf{0} \\ \gamma_{gq}^{(2)} &- \gamma_{gq}^{(2)} [\mathsf{VMV}] = \mathbf{0} \,, \gamma_{gg}^{(2)} - \gamma_{gg}^{(2)} [\mathsf{VMV}] = \mathbf{0} \end{split}$$

# Complete $N_f^2$ contributions to four-loop splitting functions in the $q \rightarrow q$ channel

#### Renormalization of $O_q$ to four loops in $q \rightarrow q$ channel

Renormalization of two-point OMEs

$$egin{aligned} &\langle q|O^{\mathsf{R}}_{q}|q
angle = Z_{qq} \left\langle q|O^{\mathsf{B}}_{q}|q
ight
angle + Z_{qg} \left\langle q|O^{\mathsf{B}}_{g}|q
ight
angle \ &+ Z_{qA} \left\langle q|O^{\mathsf{B}}_{ABC}|q
ight
angle + \left\langle q|\left[ZO\right]^{\mathsf{GV}}_{q}|q
angle, \ &Z_{qg} = \mathcal{O}(a_{s}), Z_{qA} = \mathcal{O}(a_{s}^{2}), \left[ZO\right]^{\mathsf{GV}}_{q} = \mathcal{O}(a_{s}^{3}). \end{aligned}$$

•  $\langle q | [ZO]_q^{\rm GV} | q \rangle$  vanishes at the four-loop order

- Only  $\langle q | [ZO]_q^{\text{GV}, (4)} | q 
  angle^{(0)}$  and  $\langle q | [ZO]_q^{\text{GV}, (3)} | q 
  angle^{(1)}$  are relevant
- ▶ Other operators (O<sub>q</sub>, O<sub>g</sub>, O<sub>A</sub>, O<sub>B</sub>) give all possible Lorentz structures of qq, gg, qq̄g vertex Feynman rules

$$\blacktriangleright \ \rightarrow \langle q | \left[ ZO \right]_q^{\mathrm{GV},\,(4)} \left| q \right\rangle^{(0)} = 0, \ \langle g | \left[ ZO \right]_q^{\mathrm{GV},\,(3)} \left| g \right\rangle^{(0)} = 0,$$

$$\langle q | [ZO]_q^{\mathrm{GV}, (3)} | qg \rangle^{(0)} = 0$$

#### Sample Feynman diagrams

• OMEs with physical operator insertions









• OMEs with GV operator or counterterm insertion



#### Computations of the four-loop two-point OMEs

- Focus on  $N_{f}^{2}$  part of  $\langle q|O_{q}^{\mathsf{B}}|q
  angle$  at the four-loop order, i.e.
  - Non-singlet:  $N_f^3 C_F, N_f^2 C_A C_F, N_f^2 C_F^2, N_f^2 (d^{abc})^2 / N_c$
  - Pure-singlet:  $N_f^3 C_F, N_f^2 C_A C_F, N_f^2 C_F^2$
- Working in Feynman gauge with  $\xi = 1$
- IBP reductions and derivation of DEs in parameter-x space
  - Syzygy + finite-field sampling + denominator guessing + function reconstruction implemented in Finred by Andreas von Manteuffel
- All DEs can be turned into canonical form by CANONICA combined with Libra
- The solutions of DEs are in terms of HPLs up to weight 7
- HPLs[Remiddi and Vermaseren,1999] → Harmonic Sums[Vermaseren 1998,Blumlein and Kurth,1998]; For example

$$H(1,1;x) = \sum_{n=1}^{\infty} x^n \left( -\frac{1}{n^2} + \frac{S(1,n)}{n} \right)$$

#### Results

- Managed to get  $N_f^2$  contributions with all-n dependence for both non-singlet and pure-singlet anomalous dimensions
- Non-singlet: HSs up to weight-5; Pure-singlet: HSs up to weight-4
- Non-singlet: cross-check against the previous results [Davies, Vogt, Ruijl, Ueda, Vermaseren, 16]
- Pure-singlet
  - ►  $N_f^3$ :

$$\gamma_{\mathsf{ps}}^{(3)}\big|_{N_{f}^{3}}-\gamma_{\mathsf{ps}}^{(3)}\big|_{N_{f}^{3}}[\mathsf{DVRUV}]=0$$

- ▶ N<sub>f</sub><sup>2</sup>: the all-n results are *New*; Agree with the previous fixed n results [Falcioni, Herzog, Moch, Vogt, 23]
- Sample fixed n results with n = 1000

 $\gamma_{\mathsf{ps}}^{(3)}\big|_{n=1000} = -0.000160574 \mathit{C_FN_f^3} - 0.000723846 \mathit{C_AC_FN_f^2} + 0.00116302 \mathit{C_FN_f^2}$ 

• Our all-n results allow to derive the small- $x_B$  limit easily

### Summary

- Renormalization of physical operators require unknown GV operators
- Developed a new framework to infer splitting functions
  - Two-point OMEs are used to extract splitting functions
  - ► Multi-point (≥ 3) OMEs are required to determine counterterm Feynman rules of the GV operators
- Applied it to derive 3-loop singlet splitting functions and recovered the well known results in the literature (first in general covariant gauge)
- Obtained for the first time the complete  $N_f^2$  contributions to four-loop pure-singlet anomalous dimensions with all-*n* dependence

Legs Loops	2	3	4	5
0		$[ZO]_g^{\mathrm{GV},(2)}$	O <sub>ABC</sub>	$O_q, O_g$
1	$[ZO]_g^{\mathrm{GV},(2)}$	$O_{ABC}$	$O_g$	
2	$O_{ABC}$	$O_g \int$		
3	$O_q, O_g$			
4	<910812>N2			

#### Decomposition of splitting functions

• The general structure of quark splitting functions,

$$P_{q_iq_k} = \delta_{ik}P^V_{qq} + P^S_{qq}, \quad P_{q_i\bar{q}_k} = \delta_{ik}P^V_{q\bar{q}} + P^S_{q\bar{q}}$$

• Non-singlet and singlet splitting functions

Non-singlet: 
$$P_{ns}^{\pm} = P_{qq}^{V} \pm P_{q\bar{q}}^{V}$$
,  $P_{ns}^{V} = P^{-} + n_{f}(P_{qq}^{S} - P_{q\bar{q}}^{S})$   
Singlet:  $P_{qq} = P^{+} + n_{f}(P_{qq}^{S} + P_{q\bar{q}}^{S})$ ,  $P_{qg}$ ,  $P_{gq}$ ,  $P_{gg}$ 

Evolution of PDFs

$$\frac{dT_i^{\pm}}{d\ln\mu} = 2P_{\mathsf{ns}}^{\pm} \otimes T_i^{\pm}, \ \frac{d\sum_{k=1}^{n_f} q_k^{-}}{d\ln\mu} = 2P_{\mathsf{ns}}^{V} \otimes \sum_{k=1}^{n_f} q_k^{-}, \ i = 3, 8, \cdots n_f^2 - 1$$
$$T_3^{\pm} = u^{\pm} - d^{\pm}, T_8^{\pm} = u^{\pm} + d^{\pm} - 2s^{\pm}, \cdots, q_k^{\pm} = q_k \pm \bar{q}_k,$$
$$\frac{d}{d\ln\mu} \begin{pmatrix} \Sigma \\ g \end{pmatrix} = \begin{pmatrix} P_{qq} & P_{qg} \\ P_{gq} & P_{gg} \end{pmatrix} \otimes \begin{pmatrix} \Sigma \\ g \end{pmatrix}, \quad \Sigma = \sum_{k=1}^{n_f} q_k^{+}$$

 $P_{ns}^s$ 

#### All-n Feynman rules for $O_C$

$$\begin{split} & \underset{p_{1},p_{1},q_{3}}{\overset{p_{1},p_{3},q_{3}}{2}} \overset{p_{1},p_{4},q_{4}}{\overset{p_{2},q_{3}}{2}} \\ & \rightarrow \frac{1}{24} \frac{1 + (-1)^{n}}{2} g_{2}^{2} \Delta^{\mu_{3}} \Delta^{\mu_{4}} \left\{ f^{a_{1}a_{1}a} f^{a_{2}a_{1}a} \left( 6 \left( -\Delta \cdot p_{4} \right)^{n-2} + 6 \left( \Delta \cdot p_{3} \right)^{n-2} \right. \\ & + 6 \left( \Delta \cdot (p_{1} + p_{3}) \right)^{n-2} + 6 \left( \Delta \cdot (p_{2} + p_{3}) \right)^{n-2} - \sum_{j_{1}=0}^{n-2} \left[ \right. \\ & + \left[ \left( -\Delta \cdot p_{3} \right)^{j_{1}} + \left( -\Delta \cdot p_{4} \right)^{j_{1}} \right] \left[ 3 \left( \Delta \cdot p_{1} \right)^{n-j_{1}-2} \right. \\ & + 3 \left( \Delta \cdot p_{2} \right)^{n-j_{1}-2} + \left( \Delta \cdot (p_{1} + p_{2}) \right)^{n-j_{1}-2} \right] \\ & + 9 \left[ \left( \Delta \cdot p_{1} \right)^{n-j_{1}-2} + \left( -\Delta \cdot p_{2} \right)^{j_{1}-j_{2}} \left( \Delta \cdot (p_{2} - p_{3}) \right)^{j_{1}} + \left( \Delta \cdot (p_{1} + p_{3}) \right)^{j_{1}} \right] \right] \\ & + 13 \sum_{j_{1}=0}^{n-2} \sum_{j_{2}=0}^{j_{1}} \left[ \left( -\Delta \cdot p_{2} \right)^{j_{1}-j_{2}} \left( \Delta \cdot p_{1} \right)^{n-j_{1}-2} \right] \left( \Delta \cdot (-p_{2} - p_{3}) \right)^{j_{2}} + \left( \Delta \cdot (p_{1} + p_{3}) \right)^{j_{2}} \right] \right] \right) \\ & + f^{a_{1}a_{2}} f^{a_{3}}a_{i_{3}}a_{i_{3}} \left( -6 \left( \Delta \cdot p_{3} \right)^{n-2} - 6 \left( \Delta \cdot (p_{2} + p_{3}) \right)^{n-2} \\ & + \sum_{j_{1}=0}^{n-2} \left[ 3 \left( -\Delta \cdot p_{4} \right)^{j_{1}} \left( \Delta \cdot p_{1} \right)^{n-j_{1}-2} \right] \\ & + 3 \left( -\Delta \cdot p_{3} \right)^{j_{1}} \left( \Delta \cdot p_{2} \right)^{n-j_{1}-2} + \left[ 5 \left( -\Delta \cdot p_{3} \right)^{j_{1}} - 4 \left( -\Delta \cdot p_{4} \right)^{j_{1}} \right] \left( \Delta \cdot (p_{1} + p_{3}) \right)^{n-j_{1}-2} \\ & + 3 \left( \left( -\Delta \cdot p_{3} \right)^{j_{1}} \left( -\Delta \cdot p_{2} \right)^{n-j_{1}-2} \right] \left( \Delta \cdot (-p_{2} - p_{3}) \right)^{j_{1}} \right] - 3 \Delta \cdot p_{2} \sum_{j_{1}=0}^{n-3} \left[ \left[ 3 \left[ \left( -\Delta \cdot p_{3} \right)^{j_{1}} \left( -\Delta \cdot p_{3} \right)^{j_{1}} \left( \Delta \cdot (p_{1} + p_{3}) \right)^{j_{2}} \right] \right] \right) \\ & + \frac{\sum_{j_{1}=0}^{n-2} \sum_{j_{1}=0}^{j_{1}} \left[ \left[ \left( -\Delta \cdot p_{3} \right)^{j_{1} - j_{2}} \left( \Delta \cdot p_{1} \right)^{n-j_{1}-2} \left( \left( \Delta \cdot (p_{1} + p_{3}) \right)^{j_{2}} - 14 \left( \Delta \cdot (p_{1} + p_{4}) \right)^{j_{2}} \right] \right] \right) \\ \\ & + \frac{\sum_{j_{1}=0}^{n-2} \sum_{j_{1}=0}^{j_{1}} \left[ \left[ \left( -\Delta \cdot p_{3} \right)^{j_{1} - j_{2}} \left( \Delta \cdot p_{1} \right)^{j_{1} + \left( -\Delta \cdot p_{4} \right)^{j_{1}} \left( \Delta \cdot (p_{1} + p_{2}) \right)^{j_{2}} \right] \right] \right) \\ \\ & + \frac{\sum_{j_{1}=0}^{n-2} \sum_{j_{1}=0}^{j_{1}} \left[ \left[ \left( -\Delta \cdot p_{2} \right)^{j_{1} - j_{2}} \left( \Delta \cdot p_{1} \right)^{j_{1} + \left( -\Delta \cdot p_{4} \right)^{j_{1} j_{1}} \left( \Delta \cdot (p_{1} + p_{4}) \right)^{j_{2}} \right] \right] \right) \right\}$$

Tong-Zhi Yang (University of Zurich) Renormalization of twist-two operators

#### Lorentz structures of a twist-two operator

- Based on the following two properties
  - A twist-two operator has spin-n and mass dimension n+2
  - Propagator-type Feynman rules like  $1/p^2$  can not appear in a vertex
- A twist-2 operator involving quarks or ghosts has one Lorentz structure only

$$\langle q|O|q+mg\rangle_{1\mathsf{Pl}}^{\mu_{1}\cdots\mu_{m},\,(0),\,(m)}=c_{m}\Delta^{\mu_{1}}\cdots\Delta^{\mu_{m}}$$

• A twist-two operator involving only gluons

- ▶ Only 1 + 3/2m(m-1) Lorentz structures for *m*-gluon Feynman rules
- m = 3:  $a_1 \Delta^{\mu_1} \Delta^{\mu_2} \Delta^{\mu_3} + a_2 \Delta^{\mu_1} \Delta^{\mu_2} \frac{p_1^{\mu_3}}{p_1^{\mu_3}} + \dots + a_{10} \Delta^{\mu_3} g^{\mu_1 \mu_2}$
- ▶ 19 for m = 4 and 31 for m = 5
- Count the mass dimension of  $a_i$ :  $[a_i] = x_i[\Delta \cdot p_j] + y_i[p_j \cdot p_k](y_i \ge 0)$   $[a_1] = n - 3 + y_1[p_j \cdot p_k] = n + 2 - 3 \rightarrow y_1 = 1$  (Linear in  $p_1^2, p_1 \cdot p_2 \cdots$ )  $[a_2] + [p_1^{\mu_3}] = n - 2 + y_2[p_j \cdot p_k] + 1 = n + 2 - 3 \rightarrow y_2 = 0$

• Why not  $a_{11} \Delta^{\mu_1} p_1^{\mu_2} p_2^{\mu_3}$ 

$$[a_{11}] + 2 + 3 \ge n - 1 + y_{11}[p_j \cdot p_k] + 2 + 3 = n + 4(\text{if } y_{11} = 0)$$

where 3 is mass dimension of the external 3 gluons. Twist-4 operators

#### Computations of single pole for one-loop multi-leg OMEs

Set all Mandelstam variables p<sub>1</sub><sup>2</sup>, p<sub>2</sub><sup>2</sup> · · · to numerical numbers and reconstruct their linear dependence



• Only two types of integrals are needed, other integrals are finite

$$I_1 = \int rac{d^d l}{i \pi^{d/2}} rac{1}{(l-q_1)^2 l^2}, \quad I_2 = \int rac{d^d l}{i \pi^{d/2}} rac{1}{(l-q_1)^2 l^2 ig(1-x \Delta \cdot (l+q_2)ig)}$$

• At most x-dependent logarithms appear in the single pole

$$I_2 = \frac{1}{\epsilon} \left[ \frac{\ln(1 - x\Delta \cdot q_1 - x\Delta \cdot q_2) - \ln(1 - x\Delta \cdot q_2)}{-x\Delta \cdot q_1} \right] + \mathcal{O}(\epsilon^0)$$

• Logarithms in x-space  $\rightarrow$  n-space

$$\ln(1 - x\Delta \cdot p_1 - x\Delta \cdot p_2) = \sum_{n=1}^{\infty} x^n \left[ \frac{-1}{n} (\Delta \cdot p_1 + \Delta \cdot p_2)^n \right]$$

• Factoring out the overall factor  $Z_{gA}^{(1)} = -\frac{C_A}{\epsilon} \frac{1}{n(n-1)}$ 

#### Computations of two-loop three-leg OMEs

• Set all Mandelstam variables  $p_1^2, p_2^2 \cdots$  to numerical numbers and

$$\Delta \cdot p_1 = 1, \ \Delta \cdot p_2 = z_1$$



- Derive DE with respect to *x*
- Difficult to solve DE in terms of special functions
- Expand DE to x<sup>100</sup> in the limit of x → 0, with the boundary conditions being two-loop three-leg integrals without operator insertions[T. G. Birthwright, E. W. N. Glover, and P. Marquard, 04]

#### Reconstruct two-loop counterterm Feynman rules

- Obtain two-loop three-leg OMEs to  $x^{96}$  or n=96
- For a fixed n, the result is a polynomial in  $z_1$
- Construct full-x or full-n results from data to n = 76 based on ansatz
- Polylogarithms to weight-3, generalized Harmonic sums to weight-2

$$G(1, 1, 1/(1+z_1); x) = \sum_{n=1}^{\infty} x^n \left[ \frac{S_1(z_1+1; n)}{n^2} + \frac{S_2(z_1+1; n)}{n} - \frac{S_{1,1}(1, z_1+1; n)}{n} - \frac{(z_1+1)^n}{n^3} \right]$$

where  $S_{1,1}\left(1, z_1+1; n\right) = \sum_{t_1=1}^n rac{1}{t_1} \sum_{t_2=1}^{t_1} rac{(1+z_1)^{t_2}}{t_2}$ 

- Due to the generalized Harmonic sums, impossible to disentangle
  - renormalization constants (no z<sub>1</sub> dependence)
  - operator Feynman rules (no high-weight ( $\geq 1$ ) functions)

A counterterm Feynman rule & infinite operator Feynman rules  $(N_2 = \infty)$