


ALL-ORDER PERTURBATIVE RESULTS FOR GAUGE FIELD THEORIES

Lorenzo Magnea

University of Torino - INFN Torino

HCTP - Edinburgh - 09/01/14



Outline

- Bugs and features of perturbation theory
- Factorization, evolution, summation
- From form factors to planar amplitudes
- Taming color exchanges
- Weaving multi-particle webs
- Outlook

BUGS AND FEATURES OF PERTURBATION THEORY




The bugs in PT

$$\mathcal{M}(Q, \alpha) = \mathcal{M}_0 \left[1 + \frac{\alpha}{\pi} C_1(Q) + \left(\frac{\alpha}{\pi} \right)^2 C_2(Q) + \dots \right]$$

The bugs in PT

$$\mathcal{M}(Q, \alpha) = \mathcal{M}_0 \left[1 + \frac{\alpha}{\pi} C_1(Q) + \left(\frac{\alpha}{\pi} \right)^2 C_2(Q) + \dots \right]$$




$$C_k = \infty_{\text{UV}} \quad \longrightarrow \quad C_k \propto \log^k \left(\frac{\Lambda}{Q} \right)$$

$$\mathcal{M} \left(\frac{Q}{\mu}, \alpha(\mu) \right) = \mathcal{M}_0 \left[1 + \frac{\alpha(\mu)}{\pi} C_1 \left(\frac{Q}{\mu} \right) + \left(\frac{\alpha(\mu)}{\pi} \right)^2 C_2 \left(\frac{Q}{\mu} \right) + \dots \right]$$

The bugs in PT

$$\mathcal{M}(Q, \alpha) = \mathcal{M}_0 \left[1 + \frac{\alpha}{\pi} C_1(Q) + \left(\frac{\alpha}{\pi} \right)^2 C_2(Q) + \dots \right]$$



$$C_k = \infty_{\text{UV}} \quad \longrightarrow \quad C_k \propto \log^k \left(\frac{\Lambda}{Q} \right)$$

$$\mathcal{M} \left(\frac{Q}{\mu}, \alpha(\mu) \right) = \mathcal{M}_0 \left[1 + \frac{\alpha(\mu)}{\pi} C_1 \left(\frac{Q}{\mu} \right) + \left(\frac{\alpha(\mu)}{\pi} \right)^2 C_2 \left(\frac{Q}{\mu} \right) + \dots \right]$$




$$C_k \left(\frac{Q}{\mu} \right) = \infty_{\text{IR}} \quad \longrightarrow \quad C_k \left(\frac{Q}{\mu} \right) \propto \log^k \left(\frac{Q}{m} \right)$$

$$\mathcal{M} \left(\frac{Q}{\mu}, \frac{Q}{\mu_f}, \alpha(\mu) \right) = \mathcal{M}_0 \left[1 + \frac{\alpha(\mu)}{\pi} C_1 \left(\frac{Q}{\mu}, \frac{Q}{\mu_f} \right) + \left(\frac{\alpha(\mu)}{\pi} \right)^2 C_2 \left(\frac{Q}{\mu}, \frac{Q}{\mu_f} \right) + \dots \right]$$

The bugs in PT

$$\mathcal{M}(Q, \alpha) = \mathcal{M}_0 \left[1 + \frac{\alpha}{\pi} C_1(Q) + \left(\frac{\alpha}{\pi} \right)^2 C_2(Q) + \dots \right]$$



$$C_k = \infty_{\text{UV}} \quad \longrightarrow \quad C_k \propto \log^k \left(\frac{\Lambda}{Q} \right)$$

$$\mathcal{M} \left(\frac{Q}{\mu}, \alpha(\mu) \right) = \mathcal{M}_0 \left[1 + \frac{\alpha(\mu)}{\pi} C_1 \left(\frac{Q}{\mu} \right) + \left(\frac{\alpha(\mu)}{\pi} \right)^2 C_2 \left(\frac{Q}{\mu} \right) + \dots \right]$$



$$C_k \left(\frac{Q}{\mu} \right) = \infty_{\text{IR}} \quad \longrightarrow \quad C_k \left(\frac{Q}{\mu} \right) \propto \log^k \left(\frac{Q}{m} \right)$$

$$\mathcal{M} \left(\frac{Q}{\mu}, \frac{Q}{\mu_f}, \alpha(\mu) \right) = \mathcal{M}_0 \left[1 + \frac{\alpha(\mu)}{\pi} C_1 \left(\frac{Q}{\mu}, \frac{Q}{\mu_f} \right) + \left(\frac{\alpha(\mu)}{\pi} \right)^2 C_2 \left(\frac{Q}{\mu}, \frac{Q}{\mu_f} \right) + \dots \right]$$



$$C_k \left(\frac{Q}{\mu}, \frac{Q}{\mu_f} \right) \propto k! \quad \longrightarrow \quad \sum_k \left(\frac{\alpha}{\pi} \right)^k C_k \rightarrow \infty$$


$$\mathcal{M}(Q, \alpha) = \mathcal{M}_{\text{pert.}}(Q, \alpha) + \mathcal{M}_{\text{non pert.}}(Q, \alpha)$$

The physics behind the bugs


An **infinite** result in an otherwise sensible theory **signals** that we have made a **mistake**

The physics behind the bugs

An **infinite** result in an otherwise sensible theory **signals** that we have made a **mistake**




- 📌 **Quantum** mechanical sum over **intermediate** states.
- 📌 Our **mistake**: control of high energy, short distances.
- 📌 **Fix**: locality, effective couplings, UV completion




The physics behind the bugs


An **infinite** result in an otherwise sensible theory **signals** that we have made a **mistake**



- 📌 **Quantum** mechanical sum over **intermediate** states.
- 📌 Our **mistake**: control of high energy, short distances.
- 📌 **Fix**: locality, effective couplings, UV completion




- 📌 **Quantum** mechanical sum over **final** states.
- 📌 Our **mistake**: wrong asymptotic states.
- 📌 **Fix**: inclusive cross sections, factorization.




The physics behind the bugs


An **infinite** result in an otherwise sensible theory **signals** that we have made a **mistake**




- Quantum mechanical sum over **intermediate** states.
- Our **mistake**: control of high energy, short distances.
- Fix**: locality, effective couplings, UV completion



- Quantum mechanical sum over **final** states.
- Our **mistake**: wrong asymptotic states.
- Fix**: inclusive cross sections, factorization.



- Vacuum state, **operator** product expansion.
- Our **mistake**: neglected operators, solutions.
- Fix**: include non-perturbative contributions.



Features, not bugs



- Quantum mechanics **does not destroy predictivity.**
- **Ultraviolet** physics can be **factorized** and **parametrized.**
- **Renormalization group** predicts asymptotic behaviors.
- **Local** effective field theories.




- We do not need exact knowledge of **asymptotic states.**
- **Infrared** physics can be **factorized** and **parametrized.**
- Infrared and collinear **logarithms** can be **resummed.**
- **Non-local** effective field theories.



- Perturbation theory **knows about its own limitations.**
- **Non-perturbative** contributions can be systematically **included.**
- **Power corrections** to observables can be **computed.**
- **Condensates, instantons, bound** states.

FACTORIZATION EVOLUTION SUMMATION



Ultraviolet factorization

All factorizations separating dynamics at different energy scales lead to resummation of logarithms of the ratio of scales.

Renormalization is a textbook example.

- Renormalization factorizes cutoff dependence.

$$G_0^{(n)}(p_i, \Lambda, g_0) = \prod_{i=1}^n Z_i^{1/2}(\Lambda/\mu, g(\mu)) G_R^{(n)}(p_i, \mu, g(\mu))$$

- Factorization requires the introduction of an arbitrarily chosen scale μ .

- Results must be independent of the arbitrary choice of μ .


$$\frac{dG_0^{(n)}}{d\mu} = 0 \quad \rightarrow \quad \frac{d \log G_R^{(n)}}{d \log \mu} = - \sum_{i=1}^n \gamma_i(g(\mu)) .$$

- The simple functional dependence of the factors is dictated by separation of variables.

- Proving factorization is the difficult step: it requires all-order diagrammatic analyses. Evolution equations follow automatically.

- Solving RG evolution resums logarithms of Q^2/μ^2 into $\alpha_s(\mu^2)$.


Infrared factorization



A gauge theory Feynman diagram with potential soft and collinear enhancements

- **Divergences** arise in **scattering** amplitudes from **leading regions** in loop momentum space.
- For **renormalized massless** theories only **soft** and **collinear** regions give divergences.
- **Soft** and **collinear** emissions have **universal** features, common to all **hard** processes.
- **Singular** contributions can be studied to **all orders** in perturbation theory.
- **Ward identities** and **power counting** lead to **decoupling** of soft, collinear and hard factors.
- A **soft-collinear factorization** theorem for **multi-particle** matrix elements follows.

Soft-collinear factorization: pictorial




A pictorial representation of soft-collinear factorization for fixed-angle scattering amplitudes

Color flow

In order to understand the **matrix structure** of the **soft function** it is sufficient to consider the simple case of **quark-antiquark** scattering.

At tree level



Tree-level diagrams and color flows for quark-antiquark scattering

For this process only **two color structures** are possible. A **basis** in the space of available color tensors is

$$c_{abcd}^{(1)} = \delta_{ab}\delta_{cd}, \quad c_{abcd}^{(2)} = \delta_{ac}\delta_{bd}$$

The **matrix element** is a **vector** in this space, and the Born cross section is

$$\mathcal{M}_{abcd} = \mathcal{M}_1 c_{abcd}^{(1)} + \mathcal{M}_2 c_{abcd}^{(2)} \longrightarrow \sum_{\text{color}} |\mathcal{M}|^2 = \sum_{J,L} \mathcal{M}_J \mathcal{M}_L^* \text{tr} \left[c_{abcd}^{(J)} \left(c_{abcd}^{(L)} \right)^\dagger \right] \equiv \text{Tr} [HS]_0$$

A virtual **soft gluon** will **reshuffle** color and mix the components of this vector

$$\text{QED : } \mathcal{M}_{\text{div}} = S_{\text{div}} \mathcal{M}_{\text{Born}} ; \quad \text{QCD : } [\mathcal{M}_{\text{div}}]_J = [S_{\text{div}}]_{JL} [\mathcal{M}_{\text{Born}}]_L$$


Operator Definitions

The precise **functional form** of this graphical factorization is

$$\mathcal{M}_L(p_i/\mu, \alpha_s(\mu^2), \epsilon) = \mathcal{S}_{LK}(\beta_i \cdot \beta_j, \alpha_s(\mu^2), \epsilon) H_K\left(\frac{p_i \cdot p_j}{\mu^2}, \frac{(p_i \cdot n_i)^2}{n_i^2 \mu^2}, \alpha_s(\mu^2)\right) \\ \times \prod_{i=1}^n \left[J_i\left(\frac{(p_i \cdot n_i)^2}{n_i^2 \mu^2}, \alpha_s(\mu^2), \epsilon\right) / \mathcal{J}_i\left(\frac{(\beta_i \cdot n_i)^2}{n_i^2}, \alpha_s(\mu^2), \epsilon\right) \right],$$

Here we introduced dimensionless **four-velocities** $\beta_i^\mu = Q p_i^\mu$, $\beta_i^2 = 0$, and **factorization vectors** n_i^μ , $n_i^2 \neq 0$ to define the jets,

$$J\left(\frac{(p \cdot n)^2}{n^2 \mu^2}, \alpha_s(\mu^2), \epsilon\right) u(p) = \langle 0 | \Phi_n(\infty, 0) \psi(0) | p \rangle.$$



where Φ_n is the **Wilson line** operator along the direction n^μ ,

$$\Phi_n(\lambda_2, \lambda_1) = P \exp \left[ig \int_{\lambda_1}^{\lambda_2} d\lambda n \cdot A(\lambda n) \right].$$

Note: Wilson lines represent **fast particles**, **not recoiling** against **soft** radiation

- The vectors n^μ :
- 🔧 Ensure **gauge invariance** of the jets.
 - 🔧 **Separate** collinear gluons from wide-angle soft ones.
 - 🔧 **Replace** other hard partons with a **collinear-safe** absorber.


Soft Matrices

The **soft function** S is a **matrix**, mixing the available color tensors. It is defined by a correlator of **Wilson lines**.

$$(c_L)_{\{a_k\}} \mathcal{S}_{LK}(\beta_i \cdot \beta_j, \epsilon) = \langle 0 | \prod_{k=1}^n [\Phi_{\beta_k}(\infty, 0)]_{a_k}^{b_k} | 0 \rangle (c_K)_{\{b_k\}} ,$$

The soft function S obeys a **matrix** RG evolution equation

$$\mu \frac{d}{d\mu} \mathcal{S}_{LK}(\beta_i \cdot \beta_j, \epsilon) = -\mathcal{S}_{LJ}(\beta_i \cdot \beta_j, \epsilon) \Gamma_{JK}^S(\beta_i \cdot \beta_j, \epsilon)$$



NOTE: Γ^S is **singular** for **massless** theories, due to overlapping **UV** and **collinear** poles.


S is a **pure counterterm**. In dimensional regularization, using $\alpha_s(\mu^2 = 0, \epsilon < 0) = 0$,

$$\mathcal{S}(\beta_i \cdot \beta_j, \alpha_s(\mu^2), \epsilon) = P \exp \left[-\frac{1}{2} \int_0^{\mu^2} \frac{d\xi^2}{\xi^2} \Gamma^S(\beta_i \cdot \beta_j, \alpha_s(\xi^2, \epsilon), \epsilon) \right] .$$

The determination of the **soft anomalous dimension matrix** Γ^S is the **keystone** of the resummation program for multiparton **amplitudes** and **cross sections**.

- 🔧 It **governs** the interplay of **color** exchange with **kinematics** in multiparton processes.
- 🔧 It is the only **source** of multiparton **correlations** for singular contributions.
- 🔧 **Collinear** effects are '**color singlet**' and can be extracted from **two-parton** scatterings.

FROM FORM FACTORS TO PLANAR AMPLITUDES




Gauge theory form factors

Form factors are matrix elements of **conserved currents**. For example for a massless Dirac fermion

$$\Gamma_\mu(p_1, p_2; \mu^2, \epsilon) \equiv \langle 0 | J_\mu(0) | p_1, p_2 \rangle = \bar{v}(p_2) \gamma_\mu u(p_1) \Gamma \left(\frac{Q^2}{\mu^2}, \alpha_s(\mu^2), \epsilon \right) .$$

Form factors obey soft-collinear factorization with **trivial color structure**.



In **dimensional regularization**, the Q^2 dependence is **fully determined** by evolution (**Sterman, LM**).

$$\Gamma(Q^2, \epsilon) = \exp \left\{ \frac{1}{2} \int_0^{-Q^2} \frac{d\xi^2}{\xi^2} \left[G(\bar{\alpha}(\xi^2, \epsilon), \epsilon) - \frac{1}{2} \gamma_K(\bar{\alpha}(\xi^2, \epsilon)) \log \left(\frac{-Q^2}{\xi^2} \right) \right] \right\} .$$

Tools of the trade:

- The **d-dimensional running coupling**, satisfying and displaying an **IR free** Wilson-Fisher **fixed point** for $\epsilon < 0$.
$$\mu \frac{\partial \bar{\alpha}}{\partial \mu} = -2\epsilon \bar{\alpha} + \hat{\beta}(\bar{\alpha})$$
- The **cusp anomalous dimension** γ_K , governing the **UV** singularity of a **cusped Wilson line**. Up to three loops it is proportional to the Casimir eigenvalue of the relevant color irrep (Casimir scaling)

$$\gamma_K^{[i]}(\alpha_s) = C_2^{[i]} \hat{\gamma}_K(\alpha_s) + \mathcal{O}(\alpha_s^4)$$

- The **collinear anomalous dimension** **G**, generating **subleading** collinear poles.

Gauge theory form factors

The exponentiation is **non trivial**: only poles up to $(1/\epsilon)^{n+1}$ appear in the exponent at n loops.

- All poles are **generated by the integration** over the scale of the **d-dimensional** coupling.
- All poles beyond $(1/\epsilon)^2$ are due to the running of the **four-dimensional** coupling.

In a **conformal** gauge theory (regulated by $\epsilon < 0$) all integrations are **trivial**.

$$\log [\Gamma (Q^2, \epsilon)] = -\frac{1}{2} \sum_{n=1}^{\infty} \left(\frac{\alpha_s(Q^2)}{\pi} \right)^n e^{-i\pi n\epsilon} \left[\frac{\gamma_K^{(n)}}{2n^2\epsilon^2} + \frac{G^{(n)}(\epsilon)}{n\epsilon} \right].$$

Exact results can be derived in the conformal case (Dixon, Sterman, LM):


$$\lim_{\epsilon \rightarrow 0} \left| \frac{\Gamma(Q^2, \epsilon)}{\Gamma(-Q^2, \epsilon)} \right|^2 = \exp \left[\frac{\pi^2}{4} \gamma_K (\alpha_s) \right].$$

$$G(\alpha_s, \epsilon) = 2 B_\delta (\alpha_s) + G_{\text{eik}} (\alpha_s),$$

- The **analytic continuation** of the form factor is **governed by the cusp** anomalous dimension.
- The **collinear** anomalous dimension has a **spin-independent** part determined by a Wilson line (eikonal) form factor. **Spin** enters **only** through the **DGLAP** kernel **B**.
- These results can be checked **at strong coupling** using **AdS/CFT** (Alday, Maldacena).

Exact results for planar amplitudes

All infrared divergences of planar gauge theory amplitudes are determined by the form factors.



Wedges for planar amplitudes

- In the planar limit, gluon exchanges are confined to wedges.
- Only one color structure (single trace) survives in the planar limit.
- The soft matrix is proportional to the identity in color space.
- Note that in a conformal theory S-matrix elements do not exist ...
- Regularization breaks conformal invariance and may be expected to determine the structure of scattering amplitudes.

- Indeed, in planar $N = 4$ Super Yang-Mills theory the results for IR divergences are largely inherited by finite parts.
- Two- and three-loop results suggested the 'ABDKS' ansatz

$$\mathcal{M}_n = \exp \left\{ \sum_{k=1}^{\infty} \left(\frac{N_c \alpha_s}{2\pi} \right)^k \left[f^{(k)}(\epsilon) M_n^{(1)}(k\epsilon) + C^{(l)} + \mathcal{O}(\epsilon) \right] \right\}$$


- The ansatz holds for four- and five-point planar amplitudes: they are 'exactly solved', using a dual superconformal invariance of planar amplitudes (Korchemsky et al.).
- At $n > 5$ points, a remainder function of conformal cross ratios of momentum invariants arises: it gives the 'true' four-dimensional dynamical content of the planar theory.

Exact results for planar amplitudes

- Remarkably, in $N = 4$ SYM planar amplitudes can be computed at strong coupling, via the AdS/CFT correspondence (Alday, Maldacena).
- The logarithm of the amplitude is the area of a minimal surface in AdS space, bounded by a polygonal Wilson loop, whose sides are determined by (light-like) external momenta.
- The area can be computed with purely geometrical methods.
- For the four-point function, in dimensional regularization,

$$\mathcal{M}_4 = \exp \left[i S_{\text{div}} + \frac{\sqrt{\lambda}}{8\pi} \left(\log \frac{s}{t} \right)^2 + \tilde{C} \right]$$

$$i S_{\text{div},s} = \frac{\sqrt{\lambda}}{2\pi} \sqrt{\left(\frac{\mu^2}{-s} \right)^\epsilon} \left[-\frac{1}{\epsilon^2} - \frac{1}{\epsilon} \frac{1 - \log 2}{2} \right]$$




Polygonal Wilson loop for strong coupling

- This exactly matches the weak coupling ABDKS ansatz, and gives expression for the cusp and collinear anomalous dimensions at strong coupling.
- Integrability can be used to construct an exact equation (Beisert, Eden, Staudacher) satisfied by the (planar) cusp anomalous dimension, matching both weak and strong coupling results.
- The remainder function can also be determined at strong coupling: matching weak and strong coupling is subject of much current research.


TAMING COLOR EXCHANGES

TAMING COLOR EXCHANGES



Surprising Simplicity

- 📌 The matrix Γ_S can be computed from the **poles** of S .
- 📌 **Computations** can be performed directly **for the exponent**: relevant diagram sets are called “**webs**”.
- 📌 Γ_S appears **highly complex** at high orders.
- 📌 **g-loop** webs directly **correlate** color and kinematics of up to **g+1** Wilson lines.



A web contributing to the soft anomalous dimension matrix

The **two-loop** calculation (Aybat, Dixon, Sterman) leads to a **surprising result**: for **any number** of **light-like** eikonal lines

$$\Gamma_S^{(2)} = \frac{\kappa}{2} \Gamma_S^{(1)} \quad \kappa = \left(\frac{67}{18} - \zeta(2) \right) C_A - \frac{10}{9} T_F C_F .$$

- ➡ **No** new kinematic dependence; **no** new matrix structure.
- ➡ κ is the two-loop coefficient of $\gamma_K(\alpha_s)$, rescaled by the appropriate **quadratic Casimir**,

$$\gamma_K^{(i)}(\alpha_s) = C^{(i)} \left[2 \frac{\alpha_s}{\pi} + \kappa \left(\frac{\alpha_s}{\pi} \right)^2 + \mathcal{O}(\alpha_s^3) \right] .$$

The Dipole Formula

For **massless** partons, the soft anomalous dimension matrix obeys a set of **exact equations** that **correlate color** exchange with **kinematics**.

The **simplest solution** to these equations is a **sum over color dipoles** (Becher, Neubert; Gardi, LM, 09). It gives an **ansatz** for the all-order singularity structure of **all** multiparton fixed-angle **massless** scattering amplitudes: the **dipole formula**.

📌 All **soft** and **collinear** singularities can be **collected** in a multiplicative operator **Z**

$$\mathcal{M} \left(\frac{p_i}{\mu}, \alpha_s(\mu^2), \epsilon \right) = Z \left(\frac{p_i}{\mu_f}, \alpha_s(\mu_f^2), \epsilon \right) \mathcal{H} \left(\frac{p_i}{\mu}, \frac{\mu_f}{\mu}, \alpha_s(\mu^2), \epsilon \right),$$

📌 **Z** contains both soft singularities from **S**, and collinear ones from the jet functions. It must **satisfy** its own matrix **RG equation**

$$\frac{d}{d \ln \mu} Z \left(\frac{p_i}{\mu}, \alpha_s(\mu^2), \epsilon \right) = - Z \left(\frac{p_i}{\mu}, \alpha_s(\mu^2), \epsilon \right) \Gamma \left(\frac{p_i}{\mu}, \alpha_s(\mu^2) \right).$$

The matrix **Γ** has a surprisingly simple **dipole structure**. It reads

$$\Gamma_{\text{dip}} \left(\frac{p_i}{\mu}, \alpha_s(\mu^2) \right) = -\frac{1}{4} \hat{\gamma}_K \left(\alpha_s(\mu^2) \right) \sum_{j \neq i} \ln \left(\frac{-2 p_i \cdot p_j}{\mu^2} \right) \mathbf{T}_i \cdot \mathbf{T}_j + \sum_{i=1}^n \gamma_{J_i} \left(\alpha_s(\mu^2) \right).$$

Note that **all singularities** are again **generated by integration** over the scale of the coupling.

Features of the dipole formula

- 🔊 All known results for IR divergences of massless gauge theory amplitudes are recovered.
- 🔊 The absence of multiparton correlations implies remarkable diagrammatic cancellations.
- 🔊 The color matrix structure is fixed at one loop: path-ordering is not needed.
- 🔊 The cusp anomalous dimension plays a very special role: a universal IR coupling.

Can this be the definitive answer for IR divergences in massless non-abelian gauge theories?

► There are precisely two sources of possible corrections.

- Quadrupole correlations may enter starting at three loops: they must be tightly constrained functions of conformal cross ratios of parton momenta.

$$\Gamma\left(\frac{p_i}{\mu}, \alpha_s(\mu^2)\right) = \Gamma_{\text{dip}}\left(\frac{p_i}{\mu}, \alpha_s(\mu^2)\right) + \Delta(\rho_{ijkl}, \alpha_s(\mu^2)) \quad , \quad \rho_{ijkl} = \frac{p_i \cdot p_j \, p_k \cdot p_l}{p_i \cdot p_k \, p_j \cdot p_l}$$


- The cusp anomalous dimension may violate Casimir scaling beyond three loops.

$$\gamma_K^{(i)}(\alpha_s) = C_i \hat{\gamma}_K(\alpha_s) + \tilde{\gamma}_K^{(i)}(\alpha_s)$$

- The functional form of Δ is further constrained by: collinear limits, Bose symmetry, bounds on weights, high-energy constraints. (Becher, Neubert; Dixon, Gardi, LM, 09).
- A four-loop analysis indicates that Casimir scaling holds (Becher, Neubert, Vernazza).
- Recent evidence for non-vanishing Δ at four loops from Regge limit (Caron-Huot).

WEAVING MULTI-PARTICLE WEBS

WEAVING MULTI-PARTICLE WEBS




Infrared exponentiation

All correlators of Wilson lines, regardless of shape, resum in exponential form.

$$S_n \equiv \langle 0 | \Phi_1 \otimes \dots \otimes \Phi_n | 0 \rangle = \exp(\omega_n)$$


Diagrammatic rules exist to compute directly the logarithm of the correlators.

$$\omega_{2,\text{QED}} =$$



Only **connected** photon **subdiagrams** contribute to the logarithm.

$$\omega_{2,\text{QCD}} =$$



Only gluon **subdiagrams** which are **two-eikonal irreducible** contribute to the logarithm. They have **modified color factors**.

For **eikonal form factors**, these diagrams are called **webs** (Gatheral; Frenkel, Taylor; Sterman).

Multiparticle webs

The concept of **web** generalizes non-trivially to the case of **multiple Wilson lines**.
(Gardi, Smillie, White, et al).

A **web** is a **set of diagrams** which **differ** only by the **order** of the **gluon attachments** on each Wilson line. They are **weighted** by **modified color factors**.


Writing each diagram as the product of its natural **color** factor and a **kinematic** factor

$$D = C(D)\mathcal{F}(D)$$

a **web** W can be expressed as a **sum of diagrams** in terms of a **web mixing matrix** R

$$W = \sum_D \tilde{C}(D)\mathcal{F}(D) = \sum_{D,D'} C(D')R(D',D)\mathcal{F}(D)$$

The **non-abelian exponentiation theorem** holds: each web has the color factor of a **fully connected** gluon subdiagram (Gardi, Smillie, White).



Computing webs

Bare Wilson-line correlators **vanish** beyond tree level in **dimensional regularization**: they are given by **scale-less integrals**. We require **renormalized** correlators, which depend on the **Minkowsky angles** between the Wilson lines.

$$S_{\text{ren}}(\gamma_{ij}, \alpha_s, \epsilon) = S_{\text{bare}}(\gamma_{ij}, \alpha_s, \epsilon) Z(\gamma_{ij}, \alpha_s, \epsilon) = Z(\gamma_{ij}, \alpha_s, \epsilon), \quad \gamma_{ij} = \frac{2\beta_i \cdot \beta_j}{\sqrt{\beta_i^2 \beta_j^2}}$$

To compute the **counterterm** **Z** we make use of an **auxiliary, IR-regularized** correlator

$$\begin{aligned} \hat{S}_{\text{ren}}(\gamma_{ij}, \alpha_s, \epsilon, m) &= \hat{S}_{\text{bare}}(\gamma_{ij}, \alpha_s, \epsilon, m) Z(\gamma_{ij}, \alpha_s, \epsilon) \\ &\equiv \exp(\omega) \exp(\zeta) = \exp\left\{\omega + \zeta + \frac{1}{2}[\omega, \zeta] + \dots\right\} \end{aligned}$$

The expression of **Z** in terms of the **anomalous dimension** **Γ** follows from **RG** arguments

$$Z = \exp\left[\frac{\alpha_s}{\pi} \frac{1}{2\epsilon} \Gamma^{(1)} + \left(\frac{\alpha_s}{\pi}\right)^2 \left(\frac{1}{4\epsilon} \Gamma^{(2)} - \frac{b_0}{4\epsilon^2} \Gamma^{(1)}\right) + \left(\frac{\alpha_s}{\pi}\right)^3 \left(\frac{1}{6\epsilon} \Gamma^{(3)} + \frac{1}{48\epsilon^2} [\Gamma^{(1)}, \Gamma^{(2)}] + \dots\right)\right]$$


Combining informations one can **get** **Γ** directly from the **logarithm** of the **regularized S**

$$\begin{aligned} \Gamma^{(1)} &= -2\omega^{(1,-1)} \\ \Gamma^{(2)} &= -4\omega^{(2,-1)} - 2\left[\omega^{(1,-1)}, \omega^{(1,0)}\right] \end{aligned} \quad \omega = \sum_{n=1}^{\infty} \sum_{k=-n}^{\infty} \left(\frac{\alpha_s}{\pi}\right)^n \epsilon^k \omega^{(n,k)}$$

Computing **regularized webs** is a game of **combinatorics** and **renormalization** theory.

Three-loop progress


The computation of the **three-loop** multi-particle **soft anomalous dimension** is **under way**.



(1113) web

Three-loop progress

The computation of the **three-loop** multi-particle **soft anomalous dimension** is **under way**.




(1113) web



(Gardi)


Three-loop progress

The computation of the **three-loop** multi-particle **soft anomalous dimension** is **under way**.



(1113) web


✓ (Gardi)



(1112) web


Three-loop progress

The computation of the **three-loop** multi-particle **soft anomalous dimension** is **under way**.



(1113) web

✓ (Gardi)




(1112) web

In progress


Three-loop progress

The computation of the **three-loop** multi-particle **soft anomalous dimension** is **under way**.




(1113) web

✓ (Gardi)



(1112) web


In progress



(1111) web


Three-loop progress

The computation of the **three-loop** multi-particle **soft anomalous dimension** is **under way**.




(1113) web

✓ (Gardi)



(1112) web


In progress




(1111) web



OUTLOOK



Applications: Higgs production




N^3 LL resummed cross section for Higgs production via gluon fusion at LHC

The p_T distribution for $gg \rightarrow H$ is known to NNLL and NNLO (Grazzini et al. 07, 10 Ahrens et al. 11)


- Resummation **reduces** scale uncertainty
- A subtle **polarization effect** uncovered but not implemented yet (Catani, Grazzini, 10)
- **NNLL three-loop** coefficient recently **revised** due to 'collinear anomaly'.

- The **total cross section** for $gg \rightarrow H$ is known to **N^3 LL** and **NNLO**, with **NLO EW** corrections.
- One of the **best-known** observables in the SM.
- A combined analysis (Ahrens et al. 11) gives a **3%** (th) + **8%** (pdf) + **1%** (mq) **uncertainty**.
- Ongoing **debate** on theoretical and pdf uncertainty (Baglio et al. 11).



NNLL resummed p_T distribution for Higgs production via gluon fusion at LHC

Applications: top production




NNLL top-antitop invariant mass spectrum compared to CDF data

The calculation of the **two-loop massive** anomalous dimension matrix makes it possible to perform **NNLL** resummation for **generic distributions** (Ahrens et al., 09).

- **Invariant pair mass** distribution shows remarkable **agreement** with **CDF** data.
- **Negligible** theoretical uncertainty.
- Different choices of **kinematics** and **frame** possible, vast **menu** of distributions available.

The **Tevatron** top-antitop **FB asymmetry** can be computed in QCD at **NNLL+NLO** (Ahrens et al., 09).

- **Negligible** impact on NLO result: the **solution** to the Tevatron puzzle is **not QCD higher orders**.



NNLL top-antitop FB asymmetry compared to CDF data

Summary

- 📌 We are developing an ever **deeper understanding** of the **perturbative** expansion of **gauge** field theories to **all orders**.
- 📌 Important **tools** in the infrared are **factorization** and **evolution** equations.
- 📌 **Conformal** gauge theories have **interesting** special properties.
- 📌 **Planar $N = 4$ Super Yang-Mills** theory may be exactly **solvable**.
- 📌 A simple **dipole formula** encodes **infrared singularities** for **any massless gauge theory** to a high degree of accuracy.
- 📌 Potential **corrections** to the dipole formula are **interesting**, highly **constrained**, and their study is **under way**.
- 📌 We now understand **non-abelian infrared exponentiation** for multi-particle amplitudes.
- 📌 The calculation of the **three-loop** multi-particle soft anomalous dimension is **advancing**, using **new technologies**.
- 📌 Controlling **IR singularities** leads to the **resummation** of potentially **large logarithms** in phenomenologically relevant **collider cross sections**.

THANK YOU!