

QCD (and QED) in a Euclidean finite volume

EuroPLEx Summer School 2021



Maxwell T. Hansen

August 23rd-27th

Five-day outline

☒ Warm-up and definitions

- ☒ Meaning of Euclidean
- ☒ Finite-volume set-up

☒ e^{-mL} volume effects

- ☒ Mass, matrix element in $\lambda\phi^4, g\phi^3$
- ☒ Pion mass
- ☒ LO-HVP for $(g - 2)_\mu$
- ☒ Bethe-Salpeter kernel

☒ $2 \rightarrow 2$ finite-volume (f.v.) formalism

- ☒ Scattering basics
- ☒ Derivation
- ☒ Generalizations
- ☒ Applications

☒ $(1+)\mathcal{I} \rightarrow 2$ f.v. formalism

- ☒ Derivation
- ☒ Expansions
- ☒ Applications

☒ $2 + \mathcal{I} \rightarrow 2$ f.v. formalism

- ☒ Derivation
- ☒ Testing the result
- ☒ Numerical explorations

☒ Non-local matrix elements

- ☒ Derivation
- ☒ Applications

☒ $3 \rightarrow 3$ f.v. formalism

- ☒ New complications
- ☒ Derivation ($E_n(L)$ to $\mathcal{K}_{\text{df},3}$)
- ☒ Integral equations ($\mathcal{K}_{\text{df},3}$ to $\mathcal{M}_3$)
- ☒ Testing the result
- ☒ Numerical explorations/calculations

☐ Finite-volume QED + QCD

- ☐ Different formalisms
- ☐ Basic examples

Five-day outline

☒ Warm-up and definitions

- ☒ Meaning of Euclidean
- ☒ Finite-volume set-up

☒ e^{-mL} volume effects

- ☒ Mass, matrix element in $\lambda\phi^4, g\phi^3$
- ☒ Pion mass
- ☒ LO-HVP for $(g - 2)_\mu$
- ☒ Bethe-Salpeter kernel

☒ $2 \rightarrow 2$ finite-volume (f.v.) formalism

- ☒ Scattering basics
- ☒ Derivation
- ☒ Generalizations
- ☒ Applications

☒ $(1+)\mathcal{I} \rightarrow 2$ f.v. formalism

- ☒ Derivation
- ☒ Expansions
- ☒ Applications

☒ $2 + \mathcal{I} \rightarrow 2$ f.v. formalism

- ☒ Derivation
- ☒ Testing the result
- ☒ Numerical explorations

☒ Non-local matrix elements

- ☒ Derivation
- ☒ Applications

☒ $3 \rightarrow 3$ f.v. formalism

- ☒ New complications
- ☒ Derivation ($E_n(L)$ to $\mathcal{K}_{\text{df},3}$)
- ☒ Integral equations ($\mathcal{K}_{\text{df},3}$ to $\mathcal{M}_3$)
- ☒ Testing the result
- ☒ Numerical explorations/calculations

☐ Finite-volume QED + QCD

- ☐ Different formalisms
- ☐ Basic examples

☐ Large-volume ideas

- ☐ Spectral functions
- ☐ Maiani-Testa

Five-day outline

☒ Warm-up and definitions

- ☒ Meaning of Euclidean
- ☒ Finite-volume set-up

☒ e^{-mL} volume effects

- ☒ Mass, matrix element in $\lambda\phi^4, g\phi^3$
- ☒ Pion mass
- ☒ LO-HVP for $(g-2)_\mu$
- ☒ Bethe-Salpeter kernel

☒ $2 \rightarrow 2$ finite-volume (f.v.) formalism

- ☒ Scattering basics
- ☒ Derivation
- ☒ Generalizations
- ☒ Applications

☒ $(1+)\mathcal{J} \rightarrow 2$ f.v. formalism

- ☒ Derivation
- ☒ Expansions
- ☒ Applications

☒ $2 + \mathcal{J} \rightarrow 2$ f.v. formalism

- ☒ Derivation
- ☒ Testing the result
- ☒ Numerical explorations

☒ Non-local matrix elements

- ☒ Derivation
- ☒ Applications

☒ $3 \rightarrow 3$ f.v. formalism

- ☒ New complications
- ☒ Derivation ($E_n(L)$ to $\mathcal{K}_{\text{df},3}$)
- ☒ Integral equations ($\mathcal{K}_{\text{df},3}$ to $\mathcal{M}_3$)
- ☒ Testing the result
- ☒ Numerical explorations/calculations

☐ Finite-volume QED + QCD

- ☐ Different formalisms
- ☐ Basic examples

☐ Large-volume ideas

- ☐ Spectral functions
- ☐ Maiani-Testa

Why QED + QCD?

- ❑ In our Universe, up and down quarks have different masses and electric charges
- ❑ Thus far we have been ignoring these effects

$$\langle \mathcal{O} \rangle_{[1+1+1, \alpha_{\text{QED}}=1/137]} \approx \langle \mathcal{O} \rangle_{[2+1, \alpha_{\text{QED}}=0]}$$

- ❑ This is expected to induce a percent-level systematic uncertainty

$$\alpha_{\text{QED}} \approx 0.7\% \qquad \frac{M_n - M_p}{M_n} \approx 0.1\%$$

- ❑ But many LQCD observables have reached percent level determinations!

Precision era LQCD

$$f_\pi = 130.2(8) \text{ MeV}$$

uncertainty = 0.6%

$$f_K = 155.7(0.7) \text{ MeV}$$

uncertainty = 0.5%



$$f_+(0) = 0.9698(17)$$

uncertainty = 0.2%

$$\delta_{\chi PT, \text{QED}}(\pi^- \rightarrow \ell^- \bar{\nu}) = 1.8 \%$$

$$\delta_{\chi PT, \text{QED}}(K^- \rightarrow \ell^- \bar{\nu}) = 1.1 \%$$

$$\delta_{\chi PT, \text{QED}}(K \rightarrow \pi \ell \bar{\nu}) = 0.5 \text{ to } 3 \%$$




Two basic strategies for QED+QCD

- Simulate with $\alpha_{\text{QED}}, m_u - m_d$

$$\langle \mathcal{O} \rangle = \int \mathcal{D}\Phi \mathcal{O} e^{-S_{\text{QCD}} - S_{\text{QED}} + \text{IB}}$$

simplifies observables, but signal may be suppressed



Borsanyi et al., Science 347 (2015) 1452-1455

- Expand in $\alpha_{\text{QED}}, m_u - m_d$



$$\langle \mathcal{O} \rangle = \int \mathcal{D}\Phi \mathcal{O} e^{-S_{\text{QCD}}} - \int \mathcal{D}\Phi [S_{\text{QED}} + \text{IB} \mathcal{O}] e^{-S_{\text{QCD}}} + \mathcal{O}(e^4)$$

signal is not suppressed, but observables are more complicated

QED in a box

Gauss's law prohibits a naive implementation of charged objects in a periodic volume

$$Q = \int d^3x j_0(t, \mathbf{x}) = \int d^3x \nabla \cdot \mathbf{E}(t, \mathbf{x}) = \int d\mathbf{S} \cdot \mathbf{E}(t, \mathbf{x}) = 0$$



Note: if expanding one can take infinite-volume QED
but putting QED and QCD in different spaces causes its own subtleties

- figures from A. Nicholson, GHP 2021 •

Many proposed methods

- ❑ Remove the global zero-mode of the gauge field (QED_{TL})
- ❑ Restrict the global zero-mode of the gauge field
- ❑ Remove the spatial zero-mode of the gauge field in each timeslice (QED_{L})
- ❑ Massive photon (QED_{M})
- ❑ $C^\star$ boundary conditions (QED_{C})

All equivalent if $L \rightarrow \infty$ before anything else
(before $a \rightarrow 0$, before $T \rightarrow \infty$, before $m_\gamma \rightarrow 0$, maybe before fitting)

Many proposed methods

- ☐ Remove the global zero-mode of the gauge field (QED_{TL})
- ☐ Restrict the global zero-mode of the gauge field
- ☐ Remove the spatial zero-mode of the gauge field in each timeslice (QED_{L})
- ☐ Massive photon (QED_{M})
- ☐ $C^\star$ boundary conditions (QED_{C})

All equivalent if $L \rightarrow \infty$ before anything else
(before $a \rightarrow 0$, before $T \rightarrow \infty$, before $m_\gamma \rightarrow 0$, maybe before fitting)

Don't mess with the global zero mode!

- ❑ Remove the global zero-mode of the gauge field (QED_{TL})

$$a_\mu = eL_\mu \int d^4x A_\mu(x) \stackrel{!}{=} 0$$

- ❑ Restrict the global zero-mode of the gauge field

$$-\pi < a_\mu < \pi$$

Both of these disrupt
the transfer matrix!
(i.e. the hamiltonian)



$$\int d^3\mathbf{x} \langle \psi(t, \mathbf{x}) \bar{\psi}(0) \rangle \neq \sum_{n,m} C_{nm}(L) e^{-t(E_n - E_m)} e^{-TE_m}$$



Must send $L \rightarrow \infty$ before making use of spectral decomposition

Many proposed methods

- ☐ Remove the global zero-mode of the gauge field (QED_{TL})
- ☐ Restrict the global zero-mode of the gauge field
- ☐ Remove the spatial zero-mode of the gauge field in each timeslice (QED_{L})
- ☐ Massive photon (QED_{M})
- ☐ $C^\star$ boundary conditions (QED_{C})

All equivalent if $L \rightarrow \infty$ before anything else
(before $a \rightarrow 0$, before $T \rightarrow \infty$, before $m_\gamma \rightarrow 0$, maybe before fitting)

QED_L

$$\int d^3x A_\mu(x) \stackrel{!}{=} 0$$

QED_L has a transfer matrix. It is a nonlocal prescription. Locality is a core property of QFT, it is a fundamental assumption behind

- ▶ Renormalizability by power counting
- ▶ Volume-independence of renormalization constants
- ▶ Operator product expansion
- ▶ Effective-theory description of long-distance physics
- ▶ Symanzik improvement program
- ▶ ...



QED_L has a
transfer matrix

QED_L is
non-local

Infinite-volume limit should be taken before the continuum limit.

- slide from A. Patella, Lattice 2016 plenary (not the meme though) •

QED_L

$$\int d^3x A_\mu(x) \stackrel{!}{=} 0$$

- ❑ In matrix elements of high-dimension operators one expects terms scaling as

$$\frac{1}{aL^n}$$

Vanishes if $L \rightarrow \infty$ at fixed a ,
diverges if $a \rightarrow 0$ at fixed L

- ❑ No specific evidence of a problem for quantities currently being calculated
- ❑ Failure of NREFT = important lesson
- ❑ Would be very interesting to improve understanding of these issues

Practical consequence of QED_L = modified Feynman rules in calculating volume effects

$$\frac{1}{L^3} \sum_k \longrightarrow \frac{1}{L^3} \sum_{k \neq 0}$$

Has anyone looked at exercise 3?

3.1 Evaluating the Bessel integral

In lecture we found the mass shift for $\lambda\phi^4$ theory in a periodic cubic volume:

$$\Delta m^2(L) \equiv m(L)^2 - m^2 = \frac{\lambda_0}{2} \Delta_{\mathbf{k}} \int \frac{dk_4}{2\pi} \frac{1}{k^2 + m^2},$$

where $k^2 = k_4^2 + \mathbf{k}^2$ and

$$\Delta_{\mathbf{k}} \equiv \left[\frac{1}{L^3} \sum_{\mathbf{k}} - \int \frac{d^3\mathbf{k}}{(2\pi)^3} \right].$$

3.2 Zero-mode quenching

Suppose we modify the theory such that

$$\Delta_{\mathbf{k}} \longrightarrow \Delta'_{\mathbf{k}} \equiv \left[\frac{1}{L^3} \sum_{\mathbf{k} \neq 0} - \int \frac{d^3\mathbf{k}}{(2\pi)^3} \right].$$

Give the full result for the $\mathcal{O}(\lambda)$ value of $\Delta m^2(L)$ in this case.


□ This is really $(g\phi^3)_L$

$$\Delta m(L) = \mathcal{O}(1/L^3)$$

QED_L mass determination

QED_M

- ❑ Combine Landau gauge + mass term for the photon
- ❑ Gauge invariance is broken, but in a controlled way




- Endres et al., arXiv:1507.08916, (2015) •

QED_C

- Use charge-conjugation-like boundary conditions

$$A_\mu(x + L\hat{e}_i) = -A_\mu^*(x)$$
$$\psi(x + L\hat{e}_i) = C^{-1}\bar{\psi}^T(x)$$



- Still power-like volume effects, but leading non-universal term is removed
- Exponentially suppressed flavor mixing

Not discussed here...

- ❑ Scheme dependence of separating α_{QED} from $(m_u - m_d)$ (see e.g. FLAG discussion)
- ❑ Observables beyond masses

QCD + QED captures why **lattice field theory** is so fantastic!

Methods, algorithms,
computing improves

Demand new effects
to be considered

Demand new methods and
algorithms, drive next
generation computing

Five-day outline

☒ Warm-up and definitions

- ☒ Meaning of Euclidean
- ☒ Finite-volume set-up

☒ e^{-mL} volume effects

- ☒ Mass, matrix element in $\lambda\phi^4, g\phi^3$
- ☒ Pion mass
- ☒ LO-HVP for $(g-2)_\mu$
- ☒ Bethe-Salpeter kernel

☒ $2 \rightarrow 2$ finite-volume (f.v.) formalism

- ☒ Scattering basics
- ☒ Derivation
- ☒ Generalizations
- ☒ Applications

☒ $(1+)\mathcal{I} \rightarrow 2$ f.v. formalism

- ☒ Derivation
- ☒ Expansions
- ☒ Applications

☒ $2 + \mathcal{I} \rightarrow 2$ f.v. formalism

- ☒ Derivation
- ☒ Testing the result
- ☒ Numerical explorations

☒ Non-local matrix elements

- ☒ Derivation
- ☒ Applications

☒ $3 \rightarrow 3$ f.v. formalism

- ☒ New complications
- ☒ Derivation ($E_n(L)$ to $\mathcal{K}_{\text{df},3}$)
- ☒ Integral equations ($\mathcal{K}_{\text{df},3}$ to $\mathcal{M}_3$)
- ☒ Testing the result
- ☒ Numerical explorations/calculations

☒ Finite-volume QED + QCD

- ☒ Different formalisms
- ☒ Basic examples


☐ Large-volume ideas

- ☐ Spectral functions
- ☐ Maiani-Testa


Invitation

- R-ratio based (g-2) determination

$$a^{\text{HVP,LO}}(m_\mu) = \int_{4M_\pi^2}^{\infty} ds K(m_\mu, s) R(s)$$



- Integral can be Wick rotated to Euclidean signature... but ignore this



$$\longrightarrow G(\tau) = \int_{4M_\pi^2}^{\infty} ds g(\tau, s) R(s) \qquad g(\tau, s) \propto \sqrt{s} e^{-\sqrt{s}\tau}$$

- Can we find a function $\mathcal{K}(m_\mu, \tau)$? such that...

$$\int_0^\infty d\tau \mathcal{K}(m_\mu, \tau) g(\tau, s) = K(m_\mu, s) \qquad \longrightarrow \qquad a^{\text{HVP,LO}}(m_\mu) = \int_0^\infty d\tau \mathcal{K}(m_\mu, \tau) G(\tau)$$

Of course this defines the standard kernel

General idea

we want...

$$\hat{\rho}(\bar{\omega}) \equiv \int_0^\infty d\omega \hat{\delta}_\Delta(\bar{\omega}, \omega) \rho(\omega)$$

we try to construct

$$\int_0^\infty d\tau \mathcal{K}(\bar{\omega}, \tau) e^{-\omega\tau} = \hat{\delta}_\Delta(\bar{\omega}, \omega)$$



we have...

$$G(\tau) = \int d\omega e^{-\omega\tau} \rho(\omega)$$

it then follows that


$$\hat{\rho}(\bar{\omega}) = \int_0^\infty d\tau \mathcal{K}(\bar{\omega}, \tau) G(\tau)$$

some examples...


$\hat{\delta}_\Delta(\bar{\omega}, \omega)$	$\mathcal{K}(\bar{\omega}, \tau)$
$e^{-\omega/\bar{\omega}}$ $\frac{1}{\bar{\omega} - \omega}$	$\delta(\bar{\omega} - 1/\tau)$ $e^{\bar{\omega}\tau}$
	

Role of the finite volume

$$\hat{\rho}_L(\bar{\omega}) \equiv \int_0^\infty d\omega \hat{\delta}_\Delta(\bar{\omega}, \omega) \rho_L(\omega)$$



$$G_L(\tau) = \int d\omega e^{-\omega\tau} \rho_L(\omega)$$



- Any reconstructed spectral function that $\neq$ forest of deltas...
contains implicit smearing (or else $L \rightarrow \infty$)



We require...

$$1/L \ll \Delta \ll \mu_{\text{physical}}$$

smearing function
covers many delta peaks

smearing does not overly
distort observable

R ratio



PHYSICAL REVIEW D

VOLUME 13, NUMBER 7

1 APRIL 1976

Smearing method in the quark model*

E. C. Poggio, H. R. Quinn,[†] and S. Weinberg


Lyman Laboratory of Physics, Harvard University, Cambridge, Massachusetts 02138

(Received 8 December 1975)

We propose that perturbation theory in a gauge field theory of quarks and gluons can be used to calculate physical mass-shell processes, provided that both the data and the theory are smeared over a suitable energy range. This procedure is explored in detail for the process of electron-positron annihilation into hadrons. We show that a smearing range of 3 GeV^2 in the squared center-of-mass energy should be adequate to allow the use of lowest-order perturbation theory. The smeared data are compared with theory for a variety of models. It appears to be difficult to fit the present data with theory, unless there is a new charged lepton or quark with mass in the region 2 to 3 GeV.

$$\hat{R}_\Delta(s) = \frac{\Delta}{\pi} \int_0^\infty ds' \frac{R(s')}{(s' - s)^2 + \Delta^2}$$

into the complex plane!



courtesy of M. Bruno
(using F. Jegerlehner's *alphaQED*)

Backus-Gilbert

- ❑ Developed by geophysicists Backus and Gilbert in 1967

The Resolving Power of Gross Earth Data

George Backus and Freeman Gilbert

(Received 1968 February 12)



- ❑ *Linear, model-independent* → smeared solution with a *known resolution function*

$$\begin{aligned} \sum_{\tau} \mathcal{K}(\bar{\omega}, \tau) G(\tau) &= \sum_{\tau} \mathcal{K}(\bar{\omega}, \tau) \int d\omega e^{-\omega\tau} \rho(\omega) \\ \text{BG optimizes } K &\nearrow \\ &= \int d\omega \left[\sum_{\tau} \mathcal{K}(\bar{\omega}, \tau) e^{-\omega\tau} \right] \rho(\omega) \\ &= \int d\omega \hat{\delta}_{\Delta}(\bar{\omega}, \omega) \rho(\omega) \quad \delta \text{ is exactly known} \end{aligned}$$

- ❑ The correlation matrix is used to stabilize the inverse
- ❑ The quality of the data determines the *resolution*

Backus-Gilbert (details)

□ Un-stabilized inverse

Without uncertainties, minimize the functional...

$$A[\mathcal{K}] \equiv \int_0^\infty d\omega (\bar{\omega} - \omega)^2 \hat{\delta}_\Delta(\bar{\omega}, \omega)^2 = \int_0^\infty d\omega (\bar{\omega} - \omega)^2 \left[\sum_\tau \mathcal{K}_\tau e^{-\omega\tau} \right]^2$$

with unit area constraint on δ

□ Stabilized inverse

With uncertainties, instead minimize...

$$W_\lambda[\mathcal{K}] \equiv (1 - \lambda)A[\mathcal{K}] + \lambda \frac{\mathcal{K} \cdot \text{COV} \cdot \mathcal{K}}{\mathcal{N}}$$

wants oscillating K   wants well-behaved K

λ parametrizes the tradeoff between *final resolution* and *final uncertainty*

Extended Backus Gilbert

- ❑ Important generalization from *Martin Hansen, Lupo, Tantalo*
- ❑ Addressed two limitations...

Shape and width of δ depends on data quality

No clear way to set λ

$$W_\lambda[\mathcal{K}] \equiv (1 - \lambda)A[\mathcal{K}] + \lambda \frac{\mathcal{K} \cdot \text{COV} \cdot \mathcal{K}}{\mathcal{N}}$$


Change minimizing functional

$$A[\mathcal{K}] \equiv \int_0^\infty d\omega (\bar{\omega} - \omega)^2 \hat{\delta}_\Delta(\bar{\omega}, \omega)^2$$

$$\tilde{A}[\mathcal{K}] \equiv \int_0^\infty d\omega \left[\hat{\delta}_\Delta^{\text{target}}(\bar{\omega}, \omega) - \hat{\delta}_\Delta(\bar{\omega}, \omega) \right]^2$$

Extremize W w.r.t. λ

$$W_\lambda[\mathcal{K}_{\min}(\lambda)]$$






Hansen, Lupo, Tantalo (2019)



Conservation of evil

❑ Method will always fail if one tries to...

make the resolution function too narrow




move the resolution too high in energy




Intermediate summary

- ❑ Cannot solve the inverse problem, we can get $\hat{\rho}_{L,\Delta}(\bar{\omega}) \equiv \int_0^\infty d\omega \hat{\delta}_\Delta(\bar{\omega}, \omega) \rho_L(\omega)$
- ❑ Smearing is needed anyway to *suppress volume effects*




$$1/L \ll \Delta \ll \mu_{\text{physical}}$$



- ❑ Generalized Backus-Gilbert takes $\hat{\delta}_\Delta^{\text{target}}(\bar{\omega}, \omega)$ as input

defines a continuum of
inverse-problem severity



Total rate based applications

□ Hadronic tensor, heavy flavor lifetimes

$$W_{\mu\nu}(p, q) \equiv \int d^4x \, e^{iqx} \langle \pi, p | \mathcal{J}_\mu^\dagger(x) \mathcal{J}_\nu(0) | \pi, p \rangle$$

$$\propto \int d\Phi \left| \text{diagram 1} \right|^2 + \int d\Phi \left| \text{diagram 2} \right|^2 + \int d\Phi \left| \text{diagram 3} \right|^2 + \dots$$

$$= \int_0^\infty d\omega \, e^{-\omega\tau} W_{\mu\nu}(p, q)_{\omega=p^0+q^0}$$

$$W_{\mu\nu} = \lim_{\Delta \rightarrow 0} \lim_{L \rightarrow \infty} \widehat{W}_{\mu\nu; \Delta, L}$$

□ What about *scattering* and *transition amplitudes*?

Amplitudes from spectral functions


□ First define the Euclidean correlator

$$G_L(\tau) = \langle \pi_{\mathbf{p}_4} | \pi(\tau_3, \mathbf{p}_3) \pi(0) | \pi_{\mathbf{p}_1} \rangle_L = \int_0^\infty dE_3 \rho(E_3) e^{-E_3 \tau_3}$$


and the smeared spectral function

$$\hat{\rho}_{\mathbf{p}_4 \mathbf{p}_1}^{L, \epsilon}(q_3) = \int_0^\infty dE_3 \frac{1}{q_3^0 - E_3 + i\epsilon} \rho(E_3) = \int_0^\infty dE_3 \hat{\delta}_\epsilon(q_3^0, E_2) \rho(E_3)$$

$$\frac{1}{q_3^0 - E_3 + i\epsilon} =$$



$-i$



Amplitudes from spectral functions

□ First define the Euclidean correlator

$$G_L(\tau) = \langle \pi_{\mathbf{p}_4} | \pi(\tau_3, \mathbf{p}_3) \pi(0) | \pi_{\mathbf{p}_1} \rangle_L = \int_0^\infty dE_3 \rho(E_3) e^{-E_3 \tau_3}$$

and the smeared spectral function

$$\hat{\rho}_{\mathbf{p}_4 \mathbf{p}_1}^{L, \epsilon}(q_3) = \int_0^\infty dE_3 \frac{1}{q_3^0 - E_3 + i\epsilon} \rho(E_3) = \int_0^\infty dE_3 \hat{\delta}_\epsilon(q_3^0, E_3) \rho(E_3)$$

□ Next project *on shell* at finite ϵ

$$\mathcal{M}_c^{L, \epsilon}(p_4 p_3 | p_2 p_1) \equiv \frac{2E(\mathbf{p}_3)}{Z^{1/2}(\mathbf{p}_3)} \frac{2E(\mathbf{p}_2)}{Z^{1/2}(\mathbf{p}_2)} \epsilon^2 \hat{\rho}_{\mathbf{p}_4 \mathbf{p}_1}^{L, \epsilon}(E(\mathbf{p}_3), \mathbf{p}_3)$$

□ Finally project out the *scattering amplitude*

$$\mathcal{M}_c(p_4 p_3 | p_2 p_1) = \lim_{\epsilon \rightarrow 0^+} \lim_{L \rightarrow \infty} \mathcal{M}_c^{L, \epsilon}(p_4 p_3 | p_2 p_1)$$

Some comments

❑ Derivation based in modified **LSZ** + *signature-independence* of $\rho(E)$

❑ Holds when LSZ holds

$$\langle m, \text{out} | n, \text{in} \rangle \qquad \langle m, \text{out} | \mathcal{J}(0) | n, \text{in} \rangle$$

❑ Very challenging... but systematic

for some (unknown) volume + correlator quality, we can get $D \rightarrow \pi\pi, K\bar{K}$



❑ Some nice features...

GEVP-like operator freedom

$$G_L^{[ab]}(\tau) = \langle \pi_{\mathbf{p}_4} | \pi^a(\tau_3, \mathbf{p}_3) \pi^b(0) | \pi_{\mathbf{p}_1} \rangle_L \longrightarrow \mathcal{M}^{[ab], \epsilon, L}$$

Finite range of analyticity in ϵ

Uses overlaps



Perturbative study...


Calculate in PT

$$G_L(\tau) = \langle \pi_{\mathbf{p}_4} | \pi(\tau_3, \mathbf{p}_3) \pi(0) | \pi_{\mathbf{p}_1} \rangle_L$$

Convert to this

$$\mathcal{M}_c^{L,\epsilon}(p_4 p_3 | p_2 p_1)$$

$$\text{Im } \mathcal{M}_c^{L,\epsilon}(p_4 p_3 | p_2 p_1) = \frac{\lambda^2}{2} \frac{1}{L^3} \sum_{\mathbf{k}'}^{\Lambda} \frac{1}{(2E(\mathbf{k}'))^2} \text{Im} \left\{ \frac{1}{(E_{\text{cm}} - 2E(\mathbf{k}') + i\epsilon)} \left[1 - \frac{\epsilon^2}{4E(\mathbf{k}')^2} - \frac{\epsilon(\epsilon + 2iE(\mathbf{k}'))}{E_{\text{cm}}E(\mathbf{k}')} \right] \right\}$$



Bulava, MTH (2019)

Perturbative study...


Calculate in PT

$$G_L(\tau) = \langle \pi_{\mathbf{p}_4} | \pi(\tau_3, \mathbf{p}_3) \pi(0) | \pi_{\mathbf{p}_1} \rangle_L$$

Convert to this

$$\mathcal{M}_c^{L,\epsilon}(p_4 p_3 | p_2 p_1)$$

$$\text{Im } \mathcal{M}_c^{L,\epsilon}(p_4 p_3 | p_2 p_1) = \frac{\lambda^2}{2} \frac{1}{L^3} \sum_{\mathbf{k}'}^{\Lambda} \frac{1}{(2E(\mathbf{k}'))^2} \text{Im} \left\{ \frac{1}{(E_{\text{cm}} - 2E(\mathbf{k}') + i\epsilon)} \left[1 - \frac{\epsilon^2}{4E(\mathbf{k}')^2} - \frac{\epsilon(\epsilon + 2iE(\mathbf{k}'))}{E_{\text{cm}}E(\mathbf{k}')} \right] \right\}$$



Bulava, MTH (2019)

Connection to Maiani-Testa

- Maiani and Testa considered correlators of the form

$$G_{[p]}(\tau) = \langle \pi_{-p} | \pi_p(\tau) J(0) | 0 \rangle$$

- And showed two key points

$$G_{[0]}(\tau) = \frac{\sqrt{Z_\pi}}{2M_\pi} e^{-M_\pi \tau} f(4M_\pi^2) \left[1 - a_{\pi\pi} \sqrt{\frac{m_\pi}{\pi\tau}} + O(\tau^{-3/2}) \right]$$

$G_{[p \neq 0]}(\tau)$ is plagued by un-physical (operator dependent) contributions

- Recent work with M. Bruno connects this story to spectral function amplitudes

Variations on the Maiani-Testa approach and the inverse problem

M. Bruno^a and M. T. Hansen^b

arXiv: 2012.11488



G is a smeared spectral function



□ Correlator can be viewed as a smeared spectral function

$$G_{[p]}(\tau) = \int_0^\infty d\mathcal{E} \, \rho(\mathcal{E}) \, \hat{\delta}(\mathcal{E}, \tau)$$

$$\hat{\delta}(\mathcal{E}, \tau) = \theta(\mathcal{E} - 2m_\pi) \exp[-\mathcal{E}\tau]$$


$$\rho(\mathcal{E})$$

$$q^2 = (\mathcal{E} - \omega_p)^2 - p^2$$




Shift the peak!

$$\hat{\delta}(\mathcal{E}, \tau) = \theta(\mathcal{E} - 2m_\pi) \exp[-\mathcal{E}\tau]$$



$$\hat{\delta}^\Theta(\mathcal{E}, \tau) = \Theta(\mathcal{E} - 2m_\pi, \Delta) \exp[-\mathcal{E}\tau]$$




Reaching above threshold

□ So the Maiani and Testa correlator becomes

$$G_{[p]}^{\Theta}(\tau) = \langle \pi_{-p} | \pi_p(\tau) \Theta(\hat{H} - 2\omega_p, \Delta) J(0) | 0 \rangle$$

□ Now separating the fields gives something useful above threshold

$$G_{[p]}^{\Theta}(\tau) = \frac{\sqrt{Z_{\pi}}}{2\omega_p} e^{-\omega_p \tau} \left[\Theta(0, \Delta) \operatorname{Re} f(4\omega_p^2) - 2\mathcal{J}^{(0)}(\tau, \Delta) \operatorname{Im} f(4\omega_p^2) + \dots \right]$$



Hierarchy of $\mathcal{J}^{(n)}$ provides a useful fit function

known functional forms = distinction from the work with Bulava


Constructing the Θ correlator

$$G_{[p]}^{\Theta}(\tau) = \langle \pi_{-p} | \pi_p(\tau) \Theta(\hat{H} - 2\omega_p, \Delta) J(0) | 0 \rangle$$

□ Two main methods:

□ Backus-Gilbert and HLT method

□ GEVP



$$G_{[p]}^{\Theta}(\tau) = \sum_n \Theta(E_n - 2\omega_p, \Delta) e^{-(E_n - \omega_p)\tau} \langle \pi_{-p} | \pi_p(0) | n \rangle \langle n | J(0) | 0 \rangle$$

combine finite-volume energies and matrix elements

□ Volume effects?... Suppressed as $e^{-\Delta L}$, but more investigation is needed

Five-day outline

☒ Warm-up and definitions

- ☒ Meaning of Euclidean
- ☒ Finite-volume set-up

☒ e^{-mL} volume effects

- ☒ Mass, matrix element in $\lambda\phi^4, g\phi^3$
- ☒ Pion mass
- ☒ LO-HVP for $(g - 2)_\mu$
- ☒ Bethe-Salpeter kernel

☒ $2 \rightarrow 2$ finite-volume (f.v.) formalism

- ☒ Scattering basics
- ☒ Derivation
- ☒ Generalizations
- ☒ Applications

☒ $(1+)\mathcal{J} \rightarrow 2$ f.v. formalism

- ☒ Derivation
- ☒ Expansions
- ☒ Applications

☒ $2 + \mathcal{J} \rightarrow 2$ f.v. formalism

- ☒ Derivation
- ☒ Testing the result
- ☒ Numerical explorations

☒ Non-local matrix elements

- ☒ Derivation
- ☒ Applications

☒ $3 \rightarrow 3$ f.v. formalism

- ☒ New complications
- ☒ Derivation ($E_n(L)$ to $\mathcal{K}_{\text{df},3}$)
- ☒ Integral equations ($\mathcal{K}_{\text{df},3}$ to $\mathcal{M}_3$)
- ☒ Testing the result
- ☒ Numerical explorations/calculations

☒ Finite-volume QED + QCD

- ☒ Different formalisms
- ☒ Basic examples

☒ Large-volume ideas



- ☒ Spectral functions
- ☒ Maiani-Testa

Recap!

Three essential modifications

$$\text{observable?} = \int d^N \phi e^{-S} \left[\begin{array}{l} \text{interpolator} \\ \text{for observable} \end{array} \right]$$

To proceed we have to make *three modifications*



Also... $M_{\pi, \text{lattice}} > M_{\pi, \text{our universe}}$
(but physical masses $\rightarrow$ increasingly common)



Exponentially suppressed volume effects

$$a_\mu(L) = \text{[Feynman diagrams]} + \dots$$

The Feynman diagrams represent a series of corrections to the muon magnetic moment. Each diagram consists of a muon line (wavy line) with a loop (solid line) attached. The loop is connected to the muon line at two vertices (squares). The diagrams show various topologies of the loop, including self-energy corrections and higher-order interactions.

$$\Delta a_\mu(L) = -\frac{\alpha^2}{m_\mu^2} \sum_{\mathbf{n} \neq 0} \int \frac{dp_3}{2\pi} \underbrace{\frac{e^{-|\mathbf{n}|L\sqrt{M_\pi^2+p_3^2}}}{2\pi L|\mathbf{n}|}}_{\text{volume dependence}} \int_0^\infty dx_0 \underbrace{\hat{\mathcal{K}}(m_\mu x_0)}_{\text{kernel}} \int \frac{dk_3}{2\pi} \cos(x_0 k_3) \underbrace{\text{Re } T(-k_3^2, -k_3 p_3)}_{\text{Compton amplitude}}$$

$$\underbrace{T(k^2, k \cdot p)}_{\text{reference to EFT has disappeared}} \equiv i \lim_{\mathbf{p}' \rightarrow \mathbf{p}} \sum_{q=0, \pm 1} \int d^4x e^{ikx} \langle \mathbf{p}', q | T \mathcal{J}_\rho(x) \mathcal{J}^\rho(0) | \mathbf{p}, q \rangle$$

reference to EFT has disappeared

spacelike photons only

2 → 2: Bethe Salpeter equation

□ All orders diagrammatic expansion

$$\mathcal{M} = \begin{array}{cccccc} \text{[diagram 1]} & \text{[diagram 2]} & \text{[diagram 3]} & \text{[diagram 4]} & \text{[diagram 5]} & \text{[diagram 6]} \\ \text{[diagram 7]} & \text{[diagram 8]} & \text{[diagram 9]} & \text{[diagram 10]} & \text{[diagram 11]} & \text{[diagram 12]} \\ \text{[diagram 13]} & \text{[diagram 14]} & \text{[diagram 15]} & & & \end{array}$$

$$= \text{[blue circle]} + \text{[blue circle]} \text{---} \text{[red dot]} \text{---} \text{[blue circle]} + \text{[blue circle]} \text{---} \text{[red dot]} \text{---} \text{[blue circle]} \text{---} \text{[red dot]} \text{---} \text{[blue circle]} + \dots$$

□ Construct  and  such that this is true

$$\text{[blue circle]} = \begin{array}{cccc} \text{[diagram 1]} & \text{[diagram 2]} & \text{[diagram 3]} & \text{[diagram 4]} \end{array}$$

Bethe Salpeter kernel
(2PI in the s-channel)

$$\text{[red dot on a line]} = \text{[bare line]} + \text{[diagram 1]} + \text{[diagram 2]} + \text{[diagram 3]}$$

Fully dressed propagator

2 → 2: Derivation

□ Consider the finite-volume correlator:

$$\mathcal{M}_L(P) = \text{diagram with one blob} + \text{diagram with two blobs and a loop } L \text{ with weight } 1/L^n + \text{diagram with three blobs and two loops } L + \dots \quad \square = \sum_{\mathbf{k}}$$


For two-particle energies $(2m)^2 < s < (4m)^2$, what is the L dependence?

$$\text{diagram with two blobs and a loop } L = \text{diagram with two blobs and a PV loop} + \text{diagram with two blobs and a loop } F$$

Cut projects loop to **on-shell energies**

Defines the K matrix


$$= \left[\text{diagram with one blob} + \text{diagram with two blobs and a PV loop} + \dots \right] - \left[\text{diagram with one blob} + \text{diagram with two blobs and a PV loop} + \dots \right] \text{ loop } F \left[\text{diagram with one blob} + \text{diagram with two blobs and a PV loop} + \dots \right] + \dots$$

$$= \frac{1}{\mathcal{K}(s)^{-1} + F(P, L)}$$


2 → 2: Applications

□ Single-channel case (*pions in a p-wave*)

$$\mathcal{K}(s_n)^{-1} = \rho \cot \delta(s_n) = -F(E_n, \vec{P}, L)$$



- Dudek, Edwards, Thomas in *Phys.Rev.* D87 (2013) 034505 •

$$1 + \mathcal{J} \rightarrow 2$$

□ Finite-volume matrix element

$$C_L(P) = \int_L d^4x e^{-iPx} \langle \pi | T \{ \mathcal{J}(x) \mathcal{J}(0) \} | \pi \rangle_L$$

Insert a complete set finite-volume of states

$$\xrightarrow{E \rightarrow E_n} - \frac{L^3 \langle \pi | \mathcal{J}(0) | n, L \rangle \langle n, L | \mathcal{J}(0) | \pi \rangle}{E - E_n(L)}$$


□ Infinite-volume matrix element

$$C_L(P) = C_\infty^{i\epsilon}(s) - A_{\text{in}}^{i\epsilon}(s) \frac{1}{\mathcal{M}(s) + F^{i\epsilon}(P, L)^{-1}} A_{\text{out}}^{i\epsilon}(s)$$

$$\xrightarrow{E \rightarrow E_n} - \frac{\langle \pi | \mathcal{J}(0) | \pi\pi, \text{in} \rangle \mathcal{R}(E, P, L) \langle \pi\pi, \text{out} | \mathcal{J}(0) | \pi \rangle}{E - E_n(L)}$$

$$1 + \mathcal{J} \rightarrow 2$$


Kaon decay



Implementation by RBC/UKQCD collaboration


Lellouch, Lüscher (2001) • Kim, Sachrajda, Sharpe (2005) • Christ, Kim, Yamazaki (2005)

Time-like form factors




Relevant for muon HVP contribution to muon g-2
Meyer (2011)

Resonance transition
amplitudes



Particles with spin




Agadjanov *et al.* (2014) • Briceño, MTH, Walker-Loud (2015) • Briceño, MTH (2016)

$$2 + \mathcal{J} \rightarrow 2$$



□ Fully developed formalism for multi-hadron form factors

$$\langle \pi\pi, \text{out} | \mathcal{J}_\mu | \pi\pi, \text{in} \rangle \equiv$$




□ Continuation to the pole $\rightarrow$ *resonance form factors*

□ Must carefully treat *triangle* singularities



Two types of cuts



$$\mathbf{A}'_3 \mathbf{F} \mathbf{K}_2 \mathbf{F} \mathbf{A}_3 + \mathbf{A}'_3 \mathbf{F} [\mathbf{K}_2 \mathbf{F}]^2 \mathbf{A}_3 + \mathbf{A}'_3 \mathbf{F} [\mathbf{K}_2 \mathbf{F}]^3 \mathbf{A}_3 + \dots = \mathbf{A}'_3 \mathbf{F} \frac{1}{1 - \mathbf{K}_2 \mathbf{F}} \mathbf{K}_2 \mathbf{F} \mathbf{A}_3$$

$$\mathbf{A}'_3 \mathbf{F} \mathbf{K}_2 \mathbf{G} \mathbf{K}_2 \mathbf{F} \mathbf{A}_3 + \dots$$

$$C_L - C_\infty \supset \mathbf{A}'_3 \mathbf{F}_{33} \mathbf{A}_3 \quad \mathbf{F}_{33} \equiv \frac{1}{3} \mathbf{F} + \mathbf{F} \mathbf{K}_2 \frac{1}{1 - (\mathbf{F} + \mathbf{G}) \mathbf{K}_2} \mathbf{F}$$


single $\mathbf{F}$ requires special treatment (1/3)

have not yet considered entire diagram contributions

missing contributions from *off-shellness*

missing smooth terms (short-distance parts)

Status...



☒ Identical spin-zero, no 2-to-3, no K2 poles • MTH, Sharpe (2014, 2015) •




☒ as above... but including 2-to-3 • Briceño, MTH, Sharpe (2017) •

☒ as above... but including K2 poles • Briceño, MTH, Sharpe (2018) •

☒ Non-identical, non-degenerate spin-zero $\pi\pi\pi \rightarrow \rho\pi \rightarrow \omega \rightarrow \rho\pi \rightarrow \pi\pi\pi$
 • MTH, Romero-López, Sharpe (2020) • Blanton, Sharpe (2020, 2021)

☐ Multiple three-particle channels... Spin!



$$\mathcal{M}_3 = \sum_{i,j \in \{1,2,3\}} \mathcal{M}_3^{\text{un}}(p'_i, p_j)$$



QED in a box

Gauss's law prohibits a naive implementation of charged objects in a periodic volume



$$Q = \int d^3x j_0(t, \mathbf{x}) = \int d^3x \nabla \cdot \mathbf{E}(t, \mathbf{x}) = \int d\mathbf{S} \cdot \mathbf{E}(t, \mathbf{x}) = 0$$



Note: if expanding one can take infinite-volume QED
but putting QED and QCD in different spaces causes its own subtleties

- figures from A. Nicholson, GHP 2021 •

That's all... thanks for listening!



University of Adelaide, CSSM

